



Medical Coverage Policy

Effective Date.....11/15/2019
Next Review Date..... 9/15/2020
Coverage Policy Number 0052

Genetic Testing for Hereditary and Multifactorial Conditions

Table of Contents

Overview	1
Coverage Policy	1
Laboratory Testing	2
Genetic Counseling	2
Single Gene Germline Mutation Genetic Testing.....	3
Multi-Gene Germline Mutation Genetic Testing Panels	5
Newborn Screening.....	6
General Background	6
Laboratory Testing	6
Genetic Counseling.....	6
Single Gene Germline Mutation Testing	7
Germline Mutation Genetic Testing Panels	16
Newborn Screening.....	17
Appendix A: Professional Society/Organization Recommendations/Guidelines	17
Coding/Billing Information	23
References	29

Related Coverage Resources

- [Autism Spectrum Disorders/Pervasive Developmental Disorders: Assessment and Treatment](#)
- [Genetics](#)
- [Genetic Testing Collateral File](#)
- [Infertility Services](#)
- [Recurrent Pregnancy Loss: Diagnosis and Treatment](#)

INSTRUCTIONS FOR USE

The following Coverage Policy applies to health benefit plans administered by Cigna Companies. Certain Cigna Companies and/or lines of business only provide utilization review services to clients and do not make coverage determinations. References to standard benefit plan language and coverage determinations do not apply to those clients. Coverage Policies are intended to provide guidance in interpreting certain standard benefit plans administered by Cigna Companies. Please note, the terms of a customer's particular benefit plan document [Group Service Agreement, Evidence of Coverage, Certificate of Coverage, Summary Plan Description (SPD) or similar plan document] may differ significantly from the standard benefit plans upon which these Coverage Policies are based. For example, a customer's benefit plan document may contain a specific exclusion related to a topic addressed in a Coverage Policy. In the event of a conflict, a customer's benefit plan document always supersedes the information in the Coverage Policies. In the absence of a controlling federal or state coverage mandate, benefits are ultimately determined by the terms of the applicable benefit plan document. Coverage determinations in each specific instance require consideration of 1) the terms of the applicable benefit plan document in effect on the date of service; 2) any applicable laws/regulations; 3) any relevant collateral source materials including Coverage Policies and; 4) the specific facts of the particular situation. Coverage Policies relate exclusively to the administration of health benefit plans. Coverage Policies are not recommendations for treatment and should never be used as treatment guidelines. In certain markets, delegated vendor guidelines may be used to support medical necessity and other coverage determinations.

Overview

This Coverage Policy addresses germline pathogenic or likely pathogenic variant genetic testing for hereditary and multifactorial conditions. Germline pathogenic or likely pathogenic variants are mutations occur in the egg and sperm cells; also known as the germ cells. These variants are inherited; that is, passed down in families by blood relatives.

Some conditions, such as sickle cell disease, are caused by a single germline pathogenic variant. Other conditions, such as diabetes and heart disease, are more complex. These complex conditions are referred to as multifactorial conditions. Multifactorial conditions are also inherited, but may be caused by more than one germline pathogenic or likely pathogenic variant. Environmental factors, such as nutrition, exercise, weight, smoking, drinking alcohol and medication use may influence the observable characteristics of the condition.

Types of genetic testing used to identify germline pathogenic or likely pathogenic variants that cause hereditary and multifactorial conditions include single gene pathogenic or likely pathogenic variant testing, targeted analysis and multigene sequencing panels.

Coverage Policy

Many benefit plans limit coverage of laboratory tests, genetic counseling and genetic testing. Please refer to the applicable benefit plan language to determine benefit availability and terms, conditions and limitations of coverage for the services discussed in this Coverage Policy.

If coverage for laboratory tests, genetic counseling and genetic testing is available and disease- or condition-specific criteria for genetic testing or genetic counseling are not outlined in a related Cigna Coverage Policy, the following criteria apply.

For additional information regarding coverage for specific genetic tests please refer to the [Genetic Testing Collateral File](#).

[Laboratory Testing](#)

Medically Necessary

Laboratory testing, including genetic testing (proprietary or non-proprietary, individual test or panel) is considered medically necessary when ALL of the following criteria are met:

- The proposed test or each proposed test in a panel is Food and Drug Administration (FDA)-approved and/or performed in a Clinical Laboratory Improvement Amendments (CLIA)-accredited laboratory
- The proposed test or each proposed test in a panel is medically necessary for the diagnosis(es)/indication(s) listed.
- Results of the proposed test or each proposed test in a panel will directly impact clinical decision making

For an out-of-network request to be covered at an in-network benefit level, the proposed test or each proposed test in a panel must not be available from an in-network laboratory for the indication(s) or diagnoses listed.

[Genetic Counseling](#)

Medically Necessary

Pre- and post-test genetic counseling is recommended for any individual undergoing genetic testing for any indication.

Pre-and post-test genetic counseling is considered medically necessary for EITHER of the following:

- an individual undergoing genetic testing
- an individual who is a potential candidate for genetic testing

by **ANY** of the following:

- an independent Board-Certified or Board-Eligible Medical Geneticist
- an American Board of Medical Genetics or American Board of Genetic Counseling-certified Genetic Counselor not employed by a commercial genetic testing laboratory (Genetic counselors are not excluded if they are employed by or contracted with a laboratory that is part of an Integrated Health System which routinely delivers health care services beyond just the laboratory test itself).
- a genetic nurse credentialed as either a Genetic Clinical Nurse (GCN) or an Advanced Practice Nurse in Genetics (APNG) by either the Genetic Nursing Credentialing Commission (GNCC) or the American Nurses Credentialing Center (ANCC) who is not employed by a commercial genetic testing laboratory (Genetic nurses are not excluded if they are employed by or contracted with a laboratory that is part of an Integrated Health System which routinely delivers health care services beyond just the laboratory test).

Single Gene Genetic Testing for Germline Conditions

Medically Necessary

Single gene genetic testing for a heritable disorder is considered medically necessary when **EITHER** of the following criteria is met:

- Individual demonstrates signs/symptoms of a genetically-linked heritable disease.
- Individual or fetus has a direct risk factor (e.g., based on family history or pedigree analysis) for the development of a genetically-linked heritable disease.

And **ALL** of the following criteria are met:

- results will directly impact clinical decision-making and/or clinical outcome for the individual being tested
- testing method is considered scientifically valid for identification of a genetically-linked heritable disease
- if testing guidelines exist, the clinical scenario falls within those recommendations
- the clinical benefit of testing outweighs the potential risk of psychological or medical harm to the individual being tested

Single gene genetic testing is considered medically necessary when the above criteria are met for **ALL** of the following conditions including but not limited to:

Alpha-1 antitrypsin disease	Niemann-Pick disease
Alpha and beta thalassemia	Nuclear mitochondrial genes
Canavan disease	Retinoblastoma
Cystic fibrosis	Rett syndrome
DFNB1 nonsyndromic hearing loss and deafness	Sickle cell disease
Familial hypercholesterolemia (homozygous and heterozygous)	Taya-Sachs disease
Fragile X syndrome	Von Hippel-Lindau disease
Gaucher disease	21-hydroxylase deficiency
Muscular dystrophies (DMD, BMD, EDMD, DM1, DM2, SM)	

Genetic testing with targeted mutation analysis for coagulation factor V Leiden (i.e., 1691G to A nucleotide variant) and coagulation factor II (i.e., 20210G to A nucleotide variant) is considered medically necessary for **ANY** of the following indications:

- pregnant woman who has a personal history of an unprovoked venous thromboembolism (VTE) (e.g., not associated with fracture, surgery, prolonged immobilization, cancer)
- in an individual with an unprovoked VTE when test results will impact long term medication management and at least one of the following:
 - concern for homozygous F2 or F5 or compound heterozygous F2/F5
 - high risk of recurrent VTE
- individual who has a first-degree relative* with factor V Leiden thrombophilia or F2 G2021A (prothrombin) thrombophilia and ONE of the following:
 - surgery is planned
 - pregnant
 - female who is considering estrogen contraception or hormone replacement therapy if results would influence decision to use estrogen

*A first-degree relative is defined as a blood relative with whom an individual shares approximately 50% of his/her genes, including the individual's parents, full siblings, and children.

Not Medically Necessary

Genetic testing for coagulation factor V Leiden (i.e., 1691G to A nucleotide variant) or prothrombin (i.e. 20210G to A nucleotide variant) for ANY of the following indications is considered not medically necessary (this list may not be all-inclusive):

- general population screening
- routine screening during pregnancy or prior to the use of oral contraceptives, hormone replacement therapy (HRT), or selective estrogen receptor modulators (SERMs)
- newborn testing, or routine testing in an asymptomatic child
- routine initial testing in an individual with arterial thrombosis
- testing of an asymptomatic first-degree* relative of an individual with proven symptomatic VTE and a proven coagulation factor V Leiden or prothrombin variant, for the purpose of considering primary prophylactic anticoagulation (except, as noted above, any female who is considering estrogen contraception or hormone replacement therapy if results would influence decision to use estrogen).
- neonate or child with asymptomatic central venous catheter-related thrombosis

Genetic testing is considered not medically necessary for the screening, diagnosis or management of ANY of the following conditions because there is insufficient evidence to demonstrate improved health outcomes:

- familial amyotrophic lateral sclerosis (FALS)
- genetic variants:
 - MTHFR
 - ACE
 - AGT
 - Apolipoprotein E (APOE)
 - APP
 - Presenilin 1 (PSEN1)
 - Presenilin 2 (PSEN2)
 - Prothrombin gene variant for the screening, diagnosis or management of coronary heart disease
 - Interleukin 6-174 variant
 - Chromosome 9 9p21
 - Kinesin-like protein 6
 - rs3798220 allele-lipoprotein A variant

Genetic testing or gene mapping in the general population is considered not medically necessary.

Multi-Gene Genetic Testing Panels

Medically Necessary

Genetic testing for hereditary conditions using a multigene sequencing panel is considered medically necessary when ALL of the following criteria are met:

- results will directly impact medical management of the individual being tested
- clinical presentation is consistent with a genetic etiology
- phenotype warrants testing of multiple genes and a relevant differential diagnosis list is documented
- test results may preclude the need for multiple and/or invasive procedures or tests, follow-up, or screening that would be recommended in the absence of panel testing

Genetic testing for multifactorial diseases using a multigene sequencing panel is considered medically necessary when EITHER of the following conditions is met:

- individual demonstrates signs/symptoms of a multifactorial disease
- individual has a direct risk factor (e.g., based on family history or pedigree analysis) for the development of a multifactorial disease

And ALL of the following are met:

- the panel has been shown to improve clinical outcomes
- results will directly impact clinical decision-making and clinical outcome for the individual being tested
- presence of genetic variant(s) is highly predictive for the development of the multifactorial condition

Genetic testing for nonsyndromic forms of hearing loss using a multigene sequencing panel as recommended by the American College of Medical Genetics is considered medically necessary when ALL of the following criteria are met:

- individual lacks physical findings suggestive of a known genetic syndrome
- family history is suggestive of autosomal recessive inheritance or the individual lacks a family history of prelingual hearing loss
- medical and birth histories are not suggestive of an environmental (i.e., non-genetic) cause of hearing loss, including but not limited to:
 - otitis media
 - tympanic membrane perforation
 - temporal bone fractures
 - auditory tumors
 - congenital rubella
 - congenital syphilis
 - congenital toxoplasmosis
 - congenital malformations of the inner ear
 - congenital cytomegalovirus (CMV) infection
 - prematurity
 - exposure to noise or ototoxic drugs (e.g., aminoglycosides and cyclophosphamides)

Not Medically Necessary

Genetic screening in the general population is considered not medically necessary.

Newborn Screening

Covered

Cigna covers newborn screening for genetic disorders (e.g., screening for metabolic, endocrine, hemoglobin and other disorders) performed in accordance with state mandates.

General Background

Laboratory Testing

Some general principles apply to reimbursement of all laboratory tests. The testing method being used must be scientifically validated for each indication for which the test or panel is being proposed. Due to the high complexity of genetic tests the proposed test or each proposed test in a panel must be Food and Drug Administration (FDA)-approved and/or performed in a Clinical Laboratory Improvement Amendments (CLIA)-accredited laboratory. There are several important advantages to a test being CLIA certified including the test having a higher degree of precision and performance by trained laboratory professionals. Tests performed in CLIA-accredited laboratories must meet regulatory CLIA standards. The results of each individual test or each test in a panel must be clinically useful for the diagnoses or indications for which the test is being performed. Further, outcomes must be meaningful, that is they must directly impact clinical decision making and result in improved outcomes for the individual being tested.

Genetic Counseling

Genetic counseling is defined as the process of helping individuals understand and adapt to the medical, psychological and familial indications of genetic contributions to disease. Genetic counseling services span the life cycle from preconception counseling to infertility evaluation, prenatal genetic screening and diagnosis, and include predisposition evaluation and genetic diagnosis (National Society of Genetic Counselors [NSGC]; 2010). Genetic counseling is recommended both pre-and post-genetic test to interpret family and medical histories to assess the chance of disease occurrence and recurrence, educate regarding inheritance, testing, management prevention and resources, and counsel to promote informed choices and adaptation to risk or condition (NSGC).

A variety of genetics professionals provide these services: Board-Certified or Board-Eligible Medical Geneticists, an American Board of Medical Genetics or American Board of Genetic Counseling-certified Genetic Counselor, and genetic nurses credentialed as either a Genetic Clinical Nurse (GCN) or an Advanced Practice Nurse in Genetics (APGN) by either the Genetic Nursing Credentialing Commission (GNCC) or the American Nurses Credentialing Center (ANCC). Individuals should not be employed by a commercial genetic testing laboratory, although counseling services by these individuals are not excluded if they are employed by or contracted with a laboratory that is part of an Integrated Health System which routinely delivers health care services beyond just the laboratory test itself.

Genetic Testing

Disease can result when there is an alteration or pathogenic variant in a DNA sequence which causes the cell to produce the wrong protein, or too much or too little of the correct protein. When the pathogenic variant occurs in an egg or sperm it is referred to as a germline mutation. Germline gene mutations are inherited; that is, passed down in families by blood relatives.

Some conditions, such as sickle cell disease, are caused by a single germline pathogenic variant. Other conditions, such as diabetes and heart disease, are more complex. These complex conditions are referred to as multifactorial conditions. Multifactorial conditions are also inherited, but may be caused by more than one germline pathogenic variant. The presence of a pathogenic variant(s) may increase an individual's risk of

developing one of these conditions; however, a combination of genetic and environmental factors such as nutrition, exercise, weight, smoking, drinking alcohol and medication use may influence the observable characteristics of the condition.

Genetic testing involves the analysis of human deoxyribonucleic acid (DNA), ribonucleic acid (RNA), chromosomes, proteins, and certain metabolites in order to detect alterations or changes related to an inherited disorder. Types of genetic testing used to identify germline pathogenic variant(s) that cause hereditary and multifactorial conditions include single gene testing and targeted analysis.

The National Human Genome Research Institute Task Force on Genetic Testing ([NHGRI], 1995; Holtzman, 1997) recommended the following underlying principles to ensure the safety and effectiveness of genetic tests:

- The genotypes to be detected by a genetic test must be shown by scientifically valid methods to be associated with the occurrence of a disease, independently replicated and subject to peer review.
- Analytical sensitivity and specificity of a genetic test must be determined before it is made available in clinical practice.
- Data to establish the clinical validity of genetic tests (clinical sensitivity, specificity, and predictive value) must be collected under investigative protocols. In clinical validation, the study sample must be drawn from a group of subjects representative of the population for whom the test is intended. Formal validation for each intended use of a genetic test is needed.
- Before a genetic test can be generally accepted in clinical practice, data must be collected to demonstrate the benefits and risks that accrue from both positive and negative results.

The test must also have clinical utility. Clinical utility refers to the usefulness of the test to impact health outcomes and treatment.

Genetic testing may be used to aid in diagnosis or confirmation of a disorder in a symptomatic individual (i.e., diagnostic or confirmatory testing), to predict risk of future disease in an asymptomatic individual (i.e., predictive testing), to allow reproductive planning (i.e., reproductive carrier testing, prenatal testing of a fetus, preimplantation genetic diagnosis, and newborn screening). The scope of this policy includes diagnostic and confirmatory, single or multi-gene testing for hereditary and multifactorial conditions.

Single Gene Mutation Testing for Germline Conditions

Single gene germline genetic testing is frequently performed to diagnose or confirm the presence of a disease-causing pathogenic or likely pathogenic variant and may be appropriate if an individual demonstrates characteristics of a genetically-linked heritable disease or has a direct risk factor for the development of the specific disease in question. Diagnostic testing may also be performed to help determine the course of a disease or choice of treatment. Genetic testing for a number of genetically linked heritable conditions is supported by various professional society guidelines.

Coagulation Factor V Leiden (FVL) (i.e., ~~F5 gene mutation~~ 1691G to A nucleotide variant) and Coagulation Factor II (i.e., 20210G to A nucleotide variant) (F2) Prothrombin Thrombophilia: This Coverage Policy addresses confirmatory genetic testing for FVL and F2 prothrombin (i.e., 20210G to A nucleotide variant) thrombophilia. For discussion of genetic testing for these variants in recurrent pregnancy loss please see Cigna Coverage Policy 0514 Genetic Testing for Reproductive Carrier Screening and Prenatal Diagnosis.

Factor V Leiden (FVL) variant (i.e., Arg506Gln) is the most common heritable prothrombotic risk factor in the US, with the Leiden variant accounting for 90% to 95% of activated Protein C resistance. Both heterozygotes and homozygotes are at increased risk of occurrence or recurrence of venous thromboembolism (VTE). However, clinical expression of the FVL variant is variable and many individuals with the FVL allele never develop thrombosis (GeneReviews, 2018). The risk of VTE among pregnant women who are heterozygous for FVL without a personal history of VTE or an affected first-degree relative with a thrombotic episode before age 50 years is less than 0.3%, while this risk increases to at least 10% among pregnant women with a personal or

family history of venous thromboembolism (American College of Obstetricians and Gynecologists [ACOG], 2018).

The allele 20210G to 6 (c.*97G to A) is known to be associated with coagulation factor 2 (F2) prothrombin-related thrombophilia. Individuals heterozygous for this mutation have an approximately two-to four-fold increased risk of VTE as compared to individuals without the mutation. Individuals who are homozygous for the mutation may have a more severe thrombophilia and/or an increased risk for thrombosis; however, the number of individuals who are homozygous for the variant is small and it is difficult to determine the risk of VTE in this population (American College of Medical Genetics [ACMG], 2006; Genetics Home Reference [GHR], 2018).

Clinical Utility for Genetic Testing: There is general consensus by a number of professional societies/organizations that testing for FVL and F2 is appropriate in selected individuals as an option to inform treatment. The decision to test should be based on clinical utility, that is, the likelihood that test results will impact clinical management (American College of Chest Physicians [ACCP], 2012; 2018ACMG, 2006; ACOG, 2018; Evaluation of Genomic Applications in Practice and Prevention Working Group [EGAPP], 2011; GHR; 2018; GeneReviews, 2018). Testing allows for prophylactic and/or ongoing clinical management including thromboprophylaxis and/or modification of risk factors. Persons for whom there is professional consensus regarding clinical utility for testing are:

- pregnant woman who has a personal history of an unprovoked venous thromboembolism (VTE) (e.g., not associated with fracture, surgery, prolonged immobilization, cancer)
- in an individual with an unprovoked VTE when test results will impact long term medication management and at least one of the following:
 - concern for homozygous F2 or F5 or compound heterozygous F2/F5
 - annual risk of recurrent VTE is estimated to be between 5% and 10%
- individual who has a first-degree relative* with factor V Leiden thrombophilia or F2 G2021A (prothrombin) thrombophilia and ONE of the following:
 - surgery is planned
 - pregnant
 - Female who is considering estrogen contraception or hormone replacement therapy if results would influence decision to use estrogen

Potential consequences of identifying a thrombophilic defect in a patient with venous thromboembolism (VTE) include prolonging the anticoagulant therapy beyond three-six months, or prescribing a more aggressive thromboprophylaxis in at-risk situations such as surgery, pregnancy or prolonged immobility (Roldan, 2009).

The EGAPP (2011) notes for asymptomatic family members of index cases, no prophylaxis trials have been reported; therefore, there is no direct evidence of particular benefit to family members. Potential net harm is possible if primary prophylaxis is administered to asymptomatic family members with one or more variants, because the absolute risk of an initial venous thromboembolism (VTE) event is low, and the risk of anticoagulant-induced hemorrhage is relatively high. Although technically possible, prenatal testing of a fetus and preimplantation genetic diagnosis (PGD) to determine the presence of these variants is rarely performed. Presence of the variant only increases the relative risk for thrombophilia and is not predictive of a thrombotic event (GHR, 2018).

Federal Drug Administration (FDA): The FDA has given 510K approval to several deoxyribonucleic acid (DNA)-based laboratory tests designed to test for FV Leiden (FVL) including the INFINITI System Assay (AutoGenomics, Inc., 2007, Carlsbad, CA) and the LightCycler (PCR) Instrument Version 1.2 (Roche Applied Sciences, 2003, Indianapolis, IN).

Literature Review

Targeted genetic testing to confirm diagnosis of coagulation factor V Leiden (i.e., FVL, 1691G to A nucleotide variant) and prothrombin (i.e., F2 20210G to A nucleotide variant) is appropriate in selected populations. Professional society support for testing is available in the form of published guidelines.

No randomized controlled trials have confirmed that early identification of thrombophilia affects the risk for recurrent VTE. On behalf of the AHRQ, Segal et al. (2009) published an evidence report/technology assessment of 124 studies to determine whether FVL testing alone, or in combination with prothrombin G20210A leads to improved clinical outcomes in adults with a personal history of VTE, or in adult family members of variant-positive individuals. The authors reported that there was no direct evidence in the studies that addressed this primary objective. There was moderate evidence to support the conclusion that neither harms nor benefits of testing have been demonstrated conclusively. A single study supported the hypothesis that clinicians might change management based on test results. The authors concluded that the test results have variable clinical validity for predicting VTE in these populations and have only weak clinical utility.

Marchiori et al. (2007) conducted a systematic review of prospective studies to assess the risk of recurrent VTE associated with heterozygous carriage of FVL and prothrombin G20210A (PTM) variants. The studies included a total of 3203 patients, 557 of whom were heterozygous carriers of FV Leiden (FVL). Eleven studies were included in the review, ten (seven prospective cohort studies and three randomized trials) of which examined the risk of recurrent venous thromboembolism (VTE) in heterozygous carriers of FVL. Recurrent thromboembolism occurred in 114 of the 557 heterozygous FVL carriers (20.5%) and in 382 of the 2646 non-carriers (14.4%). This data suggests that heterozygous carriers of the FVL variant may have an increased risk of recurrent VTE when compared to non-carriers.

Wu et al. (2006) performed a systematic review and meta-analysis (i.e., Thrombosis: Risk and Economic Assessment of Thrombophilia Screening [TREATS]) to establish the risk of clinical complication associated with thrombophilia in women who used oral estrogen therapy, women who were pregnant, and individuals who had undergone major orthopedic surgery. The study also measured the relative cost-effectiveness in universal and selective, history-based screening for thrombophilia. Eighty-one studies were included in the review. The highest risk of VTE in individuals who used oral contraception and hormone replacement therapy was in women with FVL (odds ratio [OR] 15.62 and 13.16 respectively). The meta-analysis also suggested that during pregnancy, women with FVL were at a significantly higher risk to develop venous thromboembolism (VTE) and to experience recurrent pregnancy loss (OR 2.06) or late pregnancy loss (2.06). The odds ratio for the association between FVL and postoperative VTE following hip or knee replacement surgery was 1.86. The authors concluded that universal thrombophilia screening is not supported by the evidence.

Professional Society/Organizations

For a summary of professional society recommendations/guidelines regarding genetic testing for coagulation Factor V Leiden and Factor II (i.e., 20210G to A nucleotide variant) please click [here](#).

The American Board of Internal Medicine's (ABIM) Foundation Choosing Wisely® Initiative (2018):

Regarding testing for thromboembolism, the American Society of Hematology notes "Don't test for thrombophilia in adult patients with venous thromboembolism (VTE) occurring in the setting of major transient risk factors (surgery, trauma or prolonged immobility)".

Use Outside of the US: A number of professional societies/organizations support genetic testing for coagulation Factor V Leiden gene variant 1691G>A) and F2 gene variant 20210G to A nucleotide change in selected subsets.

Amyotrophic Lateral Sclerosis (ALS)

No one test can provide a definitive diagnosis of ALS; a reliable biochemical abnormality shared by all patients with the disease has not been identified. Diagnosis is made by the presence of characteristic clinical features, electrodiagnostic testing (e.g., electromyography, nerve conduction velocity), and histologic findings as well as the exclusion of other conditions with related symptoms.

A number of genetic variants, including those of the superoxide dismutase 1 (SOD1), TAR DNA binding protein (TARDP or TDP-43), FUS, and C9orf72 genes have been implicated in familial ALS. About 10%–20% of all familial cases result from a specific genetic pathogenic variant of SOD1, which is inherited in an autosomal dominant manner. Greater than 100 pathogenic or likely variants mutations have been identified. Although molecular genetic testing is available for several genes associated with familial ALS, including SOD1, the presence of these pathogenic or likely pathogenic variants may not provide prognostic information; interpretation of the

significance of a mutation regarding disease severity and progression depends on the specific variant because of the wide variability in genotype/phenotype correlations. Additionally, the absence of a pathogenic or likely pathogenic variant in a family where one has not been identified is not informative as it does not rule out familial ALS caused by other pathogenic or likely pathogenic variants.

Literature Review

Several genome-wide association indicate that no definitive or common highly penetrant allele causes sporadic or familial ALS. Additionally, a number of gene variants initially thought to be causative only for familial ALS, such as SOD1, TARDBP (TDP-43) and FUS have been identified in individuals diagnosed with sporadic ALS (Belzil, 2009; GeneReviews, 2019; Wijesekera, 2009; Paubel, 2008). These data suggest the clinical utility of genetic testing for these variants is not firmly established.

There is insufficient evidence in the published, peer-reviewed scientific literature to support the clinical utility of genetic testing for the screening, diagnosis, or management of familial ALS. The identification of a pathogenic or likely pathogenic variant does not diagnose familial ALS, and does not impact treatment or health outcomes. Data are also lacking regarding the utility of prenatal or preconception carrier testing, prenatal testing of the fetus, or its use in preimplantation genetic diagnosis (PGD).

Professional Society/Organizations

For a summary of professional society recommendations/guidelines regarding genetic testing for ALS please click [here](#).

Methylenetetrahydrofolate Reductase (NAD(P)H) (MTHFR) Gene Variants

At least 40 variants in the MTHFR gene have been identified in individuals with homocystinuria (GHR, 2015). Polymorphisms in the MTHFR gene have been associated with an increased risk of homocystinuria and neural tube defects, such as anencephaly and spinal bifida, and studied as a possible risk factor for a number of other conditions such as heart disease, stroke, preeclampsia, glaucoma, cleft palate, and certain psychiatric conditions. Increased levels of homocysteine have also been associated with an increased risk of thromboembolism (GHR, 2018). Although MTHFR has been associated with increased risk of homocystinuria; genetic testing is not indicated because these variants are not associated with thromboembolism (Raffini, 2011; Dietcher, 2009).

MTHFR variants have also been associated with an increased risk of neural tube defects, such as anencephaly or spinal bifida. The 677C>T variant is the most commonly studied (GHR, 2015). This involves a change in a single deoxyribonucleic acid (DNA) nucleotide in the MTHFR gene, which produces a form of MTHFR that has reduced activity at higher temperatures (i.e., thermolabile). Individuals with the thermolabile form of the enzyme have increased blood levels of homocysteine (GHR, 2018).

Literature Review

Although there are a number of observational studies in the published peer-reviewed scientific literature regarding the association of MTHFR variants and increased risk of homocystinuria, neural tube defects and other conditions, randomized control data are limited. Evidence to demonstrate the impact of genotyping on improved health outcomes, including disease management, is also limited.

Several variants of the MTHFR gene have been associated with increased risk of developing a number of conditions; however, its role in these conditions has not been established (Genetics Home Reference [GHR], 2018; Raffini, 2011; Dietcher, 2009). There is insufficient evidence in the published peer-reviewed scientific literature to determine the clinical utility of genetic testing and its impact on net health outcomes. Professional society consensus support for MTHFR genotyping is limited. At this time the role of genetic testing for MTHFR has not been established.

Tsai et al. (2009) reported results of a longitudinal cohort analysis of participants (n=1434) of the CARDIA study. DNA was extracted from the peripheral leukocytes of blood collected from each participant. MTHFR 677C.T genotype was determined using selective amplification. The mean of serum B vitamins and tHcy concentrations and the prevalence of folate deficiency and moderate hyperhomocysteinemia were compared in 844 Caucasian and 587 African American participants before folic acid fortification (year 0 and year 7) and after fortification (year

15). Mandatory folic acid fortification as initiated by the U.S. government in 1998 improved the nutritional status of folate in both Caucasians and African Americans, with an approximate three-fold increase in folate concentrations at year 15 compared with year 0. The authors used the sensitivity and specificity of MTHFR 677C.T genotyping to predict elevated tHcy concentrations using various tHcy cutoffs to define hyperhomocysteinemia. The authors concluded that after folic acid fortification in the US, measurement of tHcy rather than genotyping of MTHFR 677TT should be used as the primary assay for the diagnosis and monitoring of moderate hyperhomocysteinemia.

Professional Societies/Organizations

For a summary of professional society recommendations/guidelines regarding genetic testing for MTHFR please click [here](#).

ACE and AGT Gene Variants

ACE Gene

The ACE gene (i.e., angiotensin I converting enzyme [peptidyl-dipeptidase A] 1) is part of the rennin-angiotensin system (GHR, 2018). ACE is a relatively nonspecific peptidase and one of the most polymorphic genes, thought to affect a number of physiologic processes including blood pressure control, hematopoiesis, reproduction, renal development, renal function, and immune response. Specifically, variants in the ACE gene have been identified as the most common cause of renal tubular dysgenesis; at least 33 variants have been found in affected individuals. A variation in the ACE gene, called the ACE I (insertion)/D (deletion) type, is a focus of ongoing research. Individuals may have two I alleles (II), two D alleles (DD), or one of each (ID). The DD type has been associated with increased levels of angiotensin-converting enzyme compared to the other types. Researchers propose that individuals with the DD type have an increased risk of stroke. It is also thought that individuals with this type who have diabetes mellitus have an increased risk of nephropathy. The contribution of other genetic and environmental influences on these risk factors is unknown (Online Mendelian Inheritance in Man [OMIM], 2018).

Angiotensinogen (AGT) Gene Variants

Individuals with an inherited variant in the AGT gene are more likely to become hypertensive and to experience more severe forms of the disease earlier in life. AGT polymorphism may be associated with increased risk of cardiovascular disease and increased responsiveness to angiotensin converting enzyme (ACE) inhibitor therapy, salt restriction and weight loss. Analysis of the gene may have potential to help individualize therapy by determining the patient's responsiveness to certain types of antihypertensive interventions. Evidence in the peer-reviewed, published scientific literature is insufficient to support the clinical utility of this testing and does not support that the detection of AGT leads to improvement of clinical outcomes in patient management.

The AGTR1 gene (i.e., angiotensin II receptor type 1 [AT1 receptor]) is also part of the rennin-angiotensin system. Like variants associated with the ACE gene, AGTR1 gene variations have also been linked to renal tubular dysgenesis. Other variants, including the 1166A>C variant have been associated with several conditions including an increased risk for the development of essential hypertension, heart disease, and nephropathy (GHR, 2018).

Although it has been suggested that the presence of ACE and AGTR1 gene variants increase risk and susceptibility for a number of conditions the influence of environmental factors and other variables on the development of these conditions is unknown. There are limited data in the published peer-reviewed scientific literature to inform improved health outcomes using the results of this testing. Established strategies for genetic testing of this gene are lacking. Genetic testing for ACE and AGTR1 gene variants remains a continued focus of research; however, the role of such testing to inform clinical practice and improve health outcomes has not yet been established.

Literature Review: Randomized controlled trial data to inform on the ability of genetic testing to improve health outcomes are lacking. Evidence in the published peer-reviewed scientific literature regarding genetic testing for ACE and AGTR1 gene variants is primarily limited to association studies and uncontrolled trials related to conditions for which increased risk has been proposed. There are scarce data regarding testing strategies and the outcomes of genetic testing on the diagnosis and management of these conditions.

Apolipoprotein E (APOE) Gene Variants

Genetic testing for apolipoprotein-E epsilon (APOE) testing has been proposed as a means to provide additional risk information for those patients currently identified as low- or intermediate-risk for cardiovascular disease by standard lipoprotein test and risk factor assessment. APOE controls the metabolism of the highly atherogenic apolipoprotein B (apo B) containing lipoproteins. It is a protein constituent of VLDL and chylomicrons. The APOE gene provides instructions for making Apo E; Apo E binds to the cell surface receptors to form molecules called lipoproteins. However, there is no uniform standard for analyzing the relationship of APOE genotypes or phenotypes to cardiovascular disease (CVD) risk. At this time genotype-phenotype correlations are incompletely understood (GeneReviews, 2014).

Genetic testing for APOE has also been proposed as a means to diagnose or predict susceptibility to early- and late-onset AD. At least three different alleles of APOE epsilon have been identified: APOE epsilon-2 (APOE e2), APOE epsilon-3 (APOE e3) and APOE epsilon-4 (APOE e4). APOE is a susceptibility polymorphism; the presence of one or two e4 alleles increases the risk but does not guarantee that someone will develop AD (Bird, 2015). Neuropathologic findings of beta-amyloid plaques and intraneuronal neurofibrillary tangles on autopsy examination remain the gold standard for diagnosis of AD (Bird, 2015). Clinical diagnosis prior to autopsy confirmation is made by use of diagnostic testing. Recommendations by the National Institute of Neurological and Communicative Diseases and Stroke and the Alzheimer's Disease and Related Disorders Association ([NINCDS-ADRDA]) criteria were published by McKhann et al. (2011), on behalf of the National Institute on Aging and the Alzheimer's Association. These criteria correctly diagnose the disease 80%-90% of the time.

The role of APOE in late-onset AD is a topic of research interest. Genotyping for the APOE e4 variant is clinically available and results in specificity estimates ranging from 75% to 81% (Genetics Home Reference [GHR], 2014; American College of Medical Genetics [ACMG], 1995; Corder, 1993). This genotype is found in many elderly persons without dementia and about 42% of persons with late-onset Alzheimer's disease (AD) do not have an apolipoprotein-E (APOE) epsilon-4 allele. The absence of this allele does not rule out the diagnosis of Alzheimer's disease, however (Mayeux, 1998). The influence of the APOE genotype on AD risk may be also modulated by cholesterol level, alpha-1 -antichymotrypsin genotype, and very low-density lipoprotein receptor gene (ACMG, 1995). According to Genetics Home Reference (GHR, 2014) association of the APOE e4 allele with AD is significant; however, APOE genotyping is neither fully specific nor sensitive. Additional genes and loci under investigation include SORL1, A2M, GSTO1 and GSTO2, GAB2, CALHM1, TOMM40, Clusterin, CR1, and PICALM (GHR, 2014; 2015; Blacker, 2002).

There is insufficient evidence in the peer-reviewed, scientific literature to support the use of APOE testing for the screening, diagnosis or management of cardiovascular disease. Neuropathological testing remains the gold standard for diagnosis of Alzheimer disease (AD). APOE genotyping does not reduce the risk of developing Alzheimer's disease, change the clinical treatment, or substantially modify disease progression in individuals with Alzheimer's disease

Literature Review

The Agency for Healthcare Research and Quality ([AHRQ], 2010) identified 15 cohort studies involving 8509 subjects that examined the association between APOE and the risk of cognitive decline. Various studies reported that APOE epsilon-4 (e4) was associated with greater decline on some, but not all, cognitive measures. Presence of an APOEe4 allele was not, however, significantly different in those who maintained cognitive performance compared to those with minor declines.

Tsuang et al. (1999) prospectively evaluated APOE testing for AD in a community-based case series of 970 persons with no previous diagnosis of dementia. Clinical diagnosis yielded a sensitivity of 84%, specificity of 50%, and positive and negative predictive values of 81% and 56%, respectively. Neuropathologic AD was confirmed in 94 of 132 patients, with a prevalence of 71%. The presence of an APOE epsilon-4 allele was associated with an estimated sensitivity of 59%, specificity of 71%, and positive and negative predictive values of 83% and 41%, respectively. The authors noted that findings do not support the use of APOE genotyping alone in the diagnosis of AD in the general medical community. In a neuropathologically confirmed series, the addition of APOE testing increased the positive predictive value of a diagnosis of AD from 90% to 94%. In those patients with a clinical diagnosis of non-Alzheimer's dementia the absence of an APOE e4 allele increased the negative predictive value from 64% to 72% (Waldemar, 2007).

Professional Societies/Organizations

For a summary of professional society recommendations/guidelines regarding genetic testing for APOE gene variants please click [here](#).

Amyloid Precursor Protein (APP), Protein Presenilin-1 (PSEN1) and Protein Presenilin 2 (PSEN2)

Early onset familial Alzheimer disease (EOFAD) is diagnosed in families with multiple affected individuals with mean age of onset before 65 years and/or with a documented pathogenic variant in one of the genes known to be associated with this disorder. Although clinically indistinguishable by phenotype, three subtypes have been recognized, based on differences in the causative gene variant: Alzheimer disease type 1 (AD1), caused by pathogenic variant of APP (10%-15% of EOFAD); Alzheimer disease type 3 (AD3), caused by pathogenic variant of PSEN1, (30%-70% of EOFAD); and Alzheimer disease type 4 (AD4), caused by pathogenic variant of PSEN2 (<5% of EOFAD). Kindreds with autosomal dominant EOFAD with no identifiable pathogenic variants in PSEN1, PSEN2, or APP have been described; thus, it is likely that variants in additional genes are causative (Bird, 2015).

Genetic testing of at-risk asymptomatic adults for early-onset familial Alzheimer's disease is clinically available for PSEN1, PSEN2 and APP variants. However, genetic testing is not helpful in predicting age of onset, severity, type of symptoms, or rate of progression in asymptomatic individuals. At this time genotyping for PSEN1, PSEN2, and APP variants does not reduce the risk of developing Alzheimer's disease, change the clinical treatment, or substantially modify disease progression in individuals with early-onset disease.

The clinical utility of genetic testing for APP, PSEN1 and PSEN2 to diagnose or manage early-onset familial Alzheimer disease in at-risk individuals or as a screening tool in the general population has not yet been established. At this time there is insufficient evidence in the published, peer-review scientific literature to demonstrate improved health outcomes with such testing.

Professional Societies/Organizations

For a summary of professional society recommendations/guidelines regarding genetic testing for APP, PSEN1 and PSEN2 gene variants please click [here](#).

Cardiac Disease-Risk Factors

Prothrombin Gene Variants: Genetic testing for prothrombin gene variants has been proposed as a means to screen, diagnose and manage coronary heart disease. Thrombosis has been implicated as a risk factor for cardiovascular disease. However, there is insufficient evidence in the published, peer-reviewed scientific literature to establish a role of genetic testing for this indication.

Interleukin 6–174 Variant: Interleukin 6 is an inflammatory cytokine that is believed to play a role in the acute phase response and inflammatory cascade similar to C-reactive protein. One polymorphism, –174, has been reported to be of specific importance (Lieb, et al., 2004). However, evidence regarding the relationship between interleukin 6–174 and cardiovascular disease has not been consistently demonstrated in the peer-reviewed, published scientific literature. The results of some studies show an association between plasma levels and cardiovascular disease (Reichert, 2016; Bermudez, et al., 2002; Ridker, et al., 2000) and, in other studies, authors have reported it is not a suitable marker for coronary heart disease and that significant associations have not been found (Sukhija, et al., 2007; Sie, et al., 2006; Lieb et al., 2004). The limitations of the overall body of published evidence preclude the ability to draw strong conclusions on the clinical utility of interleukin 6–174 testing at this time.

Kinesin-like protein 6 (KIF6): Kinesin-like protein 6 is a protein involved in intracellular transport expressed in many tissues and cell types. Theoretically, variants of KIF6 (719Arg allele) may be a risk factor associated with CVD, in particular with myocardial infarction. While the role of KIF6 in CVD is not clearly established in the peer-reviewed scientific literature, there are a few studies that support an association with CVD (Shiffman, et al., 2008a; Bare, et al., 2007; Shiffman, et al., 2008b; Iakoubova, et al., 2008). Furthermore, preliminary evidence has shown that high dose statin therapy compared with standard dose reduced the risk of death or major cardiovascular events in patients who were carriers of the gene (Iakoubova, et al., 2008). However, further studies are needed to clearly define the functional effect of the gene, the affect KIF6 has on CVD, and to determine how testing impacts medical management strategies and improves clinical outcomes.

Chromosome 9 Polymorphism 9p21: Genomic profiling (evaluating multiple genes) has recently been evaluated as a method of improving cardiac risk determination compared to traditional cardiac risk factors. The Genomic Applications in Practice and Prevention (EGAPP) Working Group (launched by the Centers for Disease Control and Prevention) sought indirect evidence to support that genomic profiling has an impact on cardiac risk estimation and that improvement in risk determination would result in management changes that improved clinical outcomes. EGAPP acknowledged direct evidence is lacking. Overall, 29 gene candidates were evaluated with 58 different gene variant associations. Only one marker, chromosome 9p21 SNPs (single nucleotide polymorphisms), had strong credibility; other combinations were moderate or weak (Palomaki, et al., 2010a). Based on the published recommendations (EGAPP, 2010) there was insufficient evidence to support testing in the general population for the 9p21 variant or for any of the 57 other variants found in 28 genes. As a result, the magnitude of health benefit for these tests were found to be insignificant. The extent to which genomic profiling alters cardiac risk estimation remains unknown and genomic testing cannot be recommended until evidence supports improved clinical outcomes.

rs3798220 allele-lipoprotein A variant: Genetic variants of the Lp(a) gene are being investigated to evaluate the influence of the variants on Lp(a) levels and associated cardiac risk. One single nucleotide polymorphism (LPA rs3798220) has been identified in the LPA gene as being associated with both elevated levels of lipoprotein(a) and an increased risk of thrombosis. Theoretically patients with a positive test for the LPA genetic variant rs3798220 may derive more benefit from the anti-thrombotic properties of aspirin due to the increased risk for thrombosis, thereby reducing cardiac disease risk. As a result, testing for the rs3798220 variant has been proposed as a method of stratifying benefit from aspirin treatment.

The U.S. Preventive Services Task Force (USPSTF) guidelines do support aspirin therapy for a specific subset of individuals for reducing the risk of stroke or myocardial infarction. Aspirin therapy is a well-established but may be associated with gastrointestinal bleeding. Authors contend that testing for the LPA genetic variant may help to better define the risk/benefit ratio of aspirin therapy when the Lp(a) level is elevated.

Evidence in the published, peer-reviewed scientific literature evaluating the association of lipoprotein A variant and elevated Lp(a) is in early stages with mixed outcomes being reported (Shiffman, et al., 2008a; Shiffman, et al., 2008b; Clarke, et al., 2009; Chasman, et al., 2009; Hopewell, et al., 2011; Anderson, et al., 2013; Koch, et al., 2013; Li, et al., 2014). Currently the evidence does not lend support that testing offers any additional prognostic value compared to Lp(a).

Gene Expression Profiling for Cardiovascular Disease Risk: Gene expression is a process by which a gene's coded information is translated into the structures present and operating in the cell and has been investigated as a diagnostic tool for evaluating individuals with cardiovascular disease.

Literature Review

Although there are some data in the published, peer-reviewed scientific literature evaluating risk factors as a method of assessing cardiovascular disease, the added value beyond that associated with traditional testing has not been firmly established. Consensus support from professional societies/organizations in the form of published guidelines is lacking. The impact of genetic testing on meaningful clinical outcomes such as morbidity and mortality has not yet been clearly defined.

Evidence in the published peer-reviewed scientific literature evaluating gene expression for determining cardiovascular disease risk (e.g., Corus CAD) is limited to prospective validation studies and case control studies (Filssoof, 2015; Ladapo, 2015; Daniels, 2014; Vargas 2013; Wingrove, et al., 2008; Rosenberg, et al., 2010; Elashoff, et al., 2011; Rosenberg, et al., 2012; Lansky, et al., 2012; McPherson, et al., 2013; Thomas, et al., 2013). Wingrove et al. (2008) and Elashoff, et al. (2011) evaluated genes associated with CAD as part of the development of the gene expression assay algorithm for assessing CAD in nondiabetic patients. Rosenberg and colleagues published results of the PREDICT trial (Personalized Risk Evaluation and Diagnosis in the Coronary Tree) in 2010, a trial designed to validate the diagnostic accuracy of gene expression, and reported sensitivity and specificity were 85% and 43% respectively. The authors noted the algorithm score was moderately correlated with maximum percent stenosis ($p < 0.001$).

As a follow-up to the 2010 trial, Rosenberg and associates reported on the relation of gene expression testing to major adverse cardiovascular events and revascularization procedures. The study group involved an extended cohort of the PREDICT trial that included the validation cohort (n=526) as well as the algorithm development cohort (n=640). Subjects underwent angiography and gene expression testing and were followed for one year post angiography (n=1116). The study endpoint was major adverse cardiac event or procedures. At one year the endpoint rate was 25% overall for all subjects. The gene expression score (GES) was associated with composite overall endpoint of both major events and procedures at one year ($p < 0.001$) and at 12 months the sensitivity and specificity were 86% and 41% respectively. Elevated GES scores (> 15) trended towards an increased rate of adverse events and procedures. The authors noted study limitations included limited follow-up period post index angiography and exclusion of individuals with high risk unstable angina and low risk asymptomatic subjects as noted by the authors. Further studies with larger cohorts and evaluation of longer term outcomes are needed.

Thomas and associates (2013) reported the results of a prospective, multicenter, double blind trial evaluating gene expression as a method to assess obstructive CAD (n=431) (COMPASS study). The study population consisted of a cohort of subjects referred for diagnostic myocardial perfusion imaging (MPI) stress testing with angina or angina equivalent symptoms. The subjects had blood samples for gene expression obtained prior to MPI and based on MPI results were referred for either invasive coronary angiography or CT angiography. The subjects were followed for 6 months with a study end point of a major adverse cardiac event. Angiography results were compared to GES and MPI results. GES was significantly correlated with maximum percent stenosis (≥ 50). Negative predictive value, sensitivity and specificity were reported at 96%, 89% and 52%, respectively. In the authors opinion gene expression scoring was more predictive of obstructive CAD compared to MPI and other clinical factors. Limitations noted by the authors included potentially lower disease prevalence in the subjects due to inclusion/exclusion criteria, and lack of comparison of GES scores to other noninvasive imaging modalities.

In another clinical trial, McPherson et al. (2013) evaluated the impact of gene expression testing on disease management by a group of cardiology specialists. The results of this study (n=88) demonstrated that subjects with low gene expression scores (i.e., ≤ 15) were more likely to have a decrease in the intensity of diagnostic testing. In addition, patients with elevated levels were more likely to undergo additional testing for the evaluation of obstructive CAD. Limitations of this study include small sample population, evaluation of short term outcomes (6 months) and inclusion criteria of low risk individuals.

Herman et al. (2013) published the results of a prospective clinical trial (n=261) to evaluate the impact of GES testing on reduction of diagnostic uncertainty in the evaluation of subjects presenting with symptoms suggestive of obstructive CAD. The trial is referred to as the "Primary Care Providers Use of a Gene Expression Test in Coronary Artery Disease Diagnosis (IMPACT-PCP)" trial. Subjects were nondiabetic patients presenting with stable, nonacute typical and atypical symptoms of obstructive CAD. Ten subjects were excluded, primarily due to GES exclusion criteria. Preliminary clinical decisions, without GES results were made by the primary care physician and compared to final decisions made with the GES results. Primary outcomes included the change in patient management between preliminary and final decisions; secondary outcomes included assessment of the pattern of change for each patient, including the effect the change had on patient outcomes. The average pretest probability of obstructive CAD was $28 \pm 17\%$. There was a change in diagnostic plan in 145 subjects with 93 having a reduction in intensity of testing ($P < .001$). GES was not associated with untoward outcomes within the first 30 days follow-up; one major adverse cardiac event occurred within the 30 day period. GES testing in this study group allowed physicians to reclassify subjects for subsequent testing. Limitations of the study included sample population of nondiabetic subjects, and short-term followup of 30 days for monitoring of adverse events.

Ladapo et al. (2014) published the results of the REGISTRY trial which was a prospective, multicenter observation registry of data collected regarding utilization of health care services for subjects at seven primary care sites who underwent GES testing. Following GES testing medical assessments of the subjects were followed for 45 days to determine how clinicians managed the subjects (e.g., cardiology referrals, cardiac stress tests, angiography). Primary outcomes included the 45 day assessment outcomes, in addition to 6 month follow up for evaluating major cardiac adverse events. The GES showed statistically significant relationships with patterns of cardiac referrals; subjects with a low GES had 94% decreased odds of referral versus subjects with an elevated GES. The overall major adverse cardiac event rate was 5/339 during the follow-up period. Ladapo and colleagues concluded GES had an effect on patient management that was clinically relevant and the test

was safe as evidenced by a low major cardiac adverse cardiac event rate. The study is limited by lack of a control group.

Professional Societies/Organizations

For a summary of professional society recommendations/guidelines regarding genetic testing for cardiac disease-risk factors please click [here](#).

Germline Genetic Testing Panels

Overall, the clinical utility of genetic testing is dependent upon the particular phenotype or observable characteristics of a disease, and set of genes being tested. Similar to genetic testing for single genes, smaller, more targeted panels to assess for a particular disorder may have clinical utility when used to aid in diagnosis of heterogeneous genetic conditions. Results of testing should directly impact clinical management and improve clinical outcomes for the individual being tested as with single gene testing. Test results may preclude the need for additional tests, follow-up or screening that would be recommended if panel testing is not performed. Additional advantages of panel testing include possible time and cost effectiveness as compared with the time and cost of analyzing each gene separately. The role of panel testing has not been established when treatment is largely supportive and/or results of testing will not result in a direct change in clinical management of the individual or lead to an improvement in clinical outcomes.

Most multi-gene panels use next-generation sequencing methodology, but some still use Sanger sequencing. Next generation sequencing technology allows large amounts DNA to be sequenced rapidly at a much lower price than prior sequencing methods. The evolution of this technology has spurred the development of tests that sequence multiple genes simultaneously. Such testing is expected to enable widespread evaluation of patients' genomes in the clinical setting (Johansen Taber, 2014). Multi-gene test panels range from small to large numbers of genes, to whole sequencing of the exome or genome.

Multi-Gene Panel Testing for Non-Syndromic Hearing Loss

More than 50% of prelingual deafness is genetic, with most of these cases being autosomal recessive and nonsyndromic. It has been noted that the reduction in the incidence of acquired causes of hearing loss has resulted in hereditary hearing loss accounting for a greater proportion of hearing loss in the general population (Hone and Smith, 2003).

Genetic forms are diagnosed by otologic, audiological, ancillary (i.e., computed tomography [CT] examination of the temporal bone), and DNA-based testing, as well as by physical examination and family history. Molecular testing of gene GJB2 (which encodes the protein connexin 26) and GJB6 should be considered in the evaluation of individuals with congenital nonsyndromic sensorineural hearing loss consistent with autosomal recessive inheritance or in families with apparent "pseudodominant" inheritance of DFNB1 (Smith, 2014). Approximately 98% of individuals with DFNB1 have two identifiable GJB2 variants. Approximately 2% of individuals with DFNB1 have one identifiable GJB2 variant and one of two large deletions that include a portion of GJB6. Diagnosis depends upon molecular genetic testing to identify deafness-causing variants in the genes GJB2 and GJB6. GJB2 gene is the major gene responsible for nonsyndromic, recessive deafness and variants in GJB2 gene and GJB6 gene together account for 50% of autosomal recessive nonsyndromic hearing loss.

Multi-gene testing or screening with a panel of genetic tests has been proposed to test for many causes of hearing loss. The extreme genetic heterogeneity and the frequent lack of phenotypic variability make genetic diagnosis of non-syndromic hearing loss (NSHL) difficult using single-gene screening techniques—for this reason, multi-gene screening panels for NSHL have been developed. For some syndromic forms of hearing loss, such as Usher syndrome or Pendred syndrome, the nonauditory features can be subtle, especially in early childhood and certain environmental or nongenetic factors play a major etiologic role in hearing loss.

Professional Societies/Organizations

For a summary of professional society recommendations/guidelines regarding genetic testing for multi-gene panel testing for non-syndromic hearing loss please click [here](#).

Newborn Screening

Newborn screening is performed to limit the morbidity and mortality attributable to selected inherited diseases (American Academy of Pediatrics ([AAP], 2013). Newborn screening programs are organized through state governments and are generally mandated. According to the March of Dimes (2015), screening is available for disorders in which accurate diagnosis and early treatment of the disorder can improve health outcomes. Some genetic screening tests are not deoxyribonucleic acid (DNA) - or chromosome-based tests but use biochemical markers or phenotypic features.

AAP guidelines (2013) note the AAP and American College of Medical Genetics (ACMG) do not support routine carrier testing in minors when such testing does not provide health benefits in childhood.

Use Outside of the US

See Appendix A for relevant statements.

Appendix A: Professional Society/Organization Recommendations/Guidelines

Factor V Leiden and Coagulation Factor II (i.e., 20210G>A variant)

American College of Chest Physicians (ACCP): On behalf of the ACCP, Guyatt et al. (2012) published The Antithrombotic Therapy and Prevention of Thrombosis, 9th ed: American College of Chest Physicians Evidence-Based Clinical Practice Guidelines that provides treatment guidelines for the circumstance when a pregnant woman is known to be homozygous for FVL or the prothrombin 20210A mutation variant

American College of Medical Genetics (ACMG): On behalf of the ACMG, Grody et al. (2006) recommended testing for the following indications:

- age under 50, any venous thrombosis
- venous thrombosis in unusual sites (such as hepatic, mesenteric, and cerebral veins)
- recurrent venous thrombosis
- venous thrombosis and a strong family history of thrombotic disease
- venous thrombosis in pregnant women or women taking oral contraceptives
- relatives of individuals with venous thrombosis under age 50
- myocardial infarction in female smokers under age 50

Testing may also be considered in the following situations:

- venous thrombosis, age over 50, except when active malignancy is present
- relatives of individuals known to have FV Leiden (FVL). Knowledge that they have the FVL variant may influence management of pregnancy and may be a factor in decision-making regarding oral contraceptive use.
- women with recurrent pregnancy loss or unexplained severe preeclampsia, placental abruption, intrauterine fetal growth retardation, or stillbirth. Knowledge of FVL carrier status may influence management of future pregnancies.

FVL testing is not recommended for the following:

- a general population screen
- a routine initial test during pregnancy or prior to the use of oral contraceptives, hormone replacement therapy (HRT) or selective estrogen receptor modulators (SERMs)
- a prenatal or newborn test, or as a routine test in asymptomatic children
- a routine initial test in individuals with arterial thrombosis (testing may be considered, however, in selected young individuals [under age 50] with unexplained arterial thrombosis in the absence of other risk factors for atherosclerotic vascular disease)

American Congress of Obstetricians and Gynecologists (ACOG): Practice Bulletin number 197 (2018) regarding inherited thrombophilias in pregnancy notes:

- Screening for inherited thrombophilias is not recommended for women with a history of fetal loss or adverse pregnancy outcomes (e.g., placental abruption, preeclampsia, fetal growth restriction) because is not recommended. (Level B, based on limited or inconsistent scientific evidence)
- Because of the lack of association between either heterozygosity or homozygosity for the MTHFR C677T polymorphism and any negative pregnancy outcomes, increased ant increased risk of VTE, screening with either MTHFR mutation analyses or fasting homocysteine levels is not recommended. (Level B, based on limited or inconsistent scientific evidence)
- Among women with personal histories of VTE, recommended screening tests for inherited thrombophilias should include factor V Leiden mutation and G20210A mutation. (Level C, based primarily on consensus and expert opinion)

Screening for inherited thrombophilias is useful only when results will affect management decisions, and is not useful in situations where treatment is indicated for other risk factors. Targeted assessment may be considered in the following clinical settings:

- A personal history of venous thromboembolism that was associated with or without a recurrent risk factor (e.g., fractures, surgery, and prolonged immobilization) and no prior thrombophilia testing.
- A first-degree relative (e.g., parent or sibling) with a history of high-risk thrombophilia.

Recommendations based primarily on consensus and expert opinion note screening for inherited thrombophilias should include FVL mutation; prothrombin G20210A mutation; and antithrombin, protein C, and protein S deficiencies. Additionally all patients with inherited thrombophilias should undergo individualized risk assessment, which may modify management decisions.

American Heart Association (AHA): On behalf of the AHA, Kim (2003) published a statement supporting the ACMG guidelines. According to this statement, patients who develop a deep vein thrombus (DVT) or pulmonary embolism (PE) and are from a family with a confirmed FVL diagnosis should be tested. Likewise, even if there is no family history of FV Leiden (FVL, F5L), anyone who has had a deep vein thrombus (DVT) or pulmonary embolus (PE) that is unexplained, recurrent, occurred at a young age (under 50), occurred during pregnancy, was associated with hormone use, or developed in an unusual site (such as the veins of the brain or abdomen) may benefit from testing for FVL and other causes of hereditary thrombophilia.

Evaluation of Genomic Applications in Practice and Prevention Working Group (EGAPP, 2011): The EGAPP published guidelines regarding the routine testing for factor V Leiden (R506Q) and prothrombin (20210G>A) mutations in adults with a history of idiopathic venous thromboembolism and their adult family members (2011). According to the EGAPP there was adequate evidence to recommend against routine testing for FVL and/or prothrombin 20210G>A (PT) in the following circumstances: (1) adults with idiopathic venous thromboembolism (VTE). In such cases, longer term secondary prophylaxis to avoid recurrence offers similar benefits to patients with and without one or more of these mutations. (2) Asymptomatic adult family members of patients with VTE and an FVL or PT mutation, for the purpose of considering primary prophylactic anticoagulation. Potential benefits are unlikely to exceed potential harms. The overall certainty of these findings was deemed "moderate." The evidence was insufficient to determine whether FVL/PT testing might have clinical utility in some circumstances, such as for identifying FVL homozygosity among asymptomatic family members of adults with idiopathic VTE or counseling patients about the risks and benefits of antithrombotic therapy. Based on the available evidence, the certainty of net health benefit was deemed "low." The recommendations do not extend to patients with other risk factors for thrombosis, such as contraceptive use, as the evidence review that serves as the basis for the recommendations focused primarily on idiopathic VTE.

Pregnancy and Thrombosis Working Group: On behalf of this association, Duhl et al. (2007) published a consensus report and recommendations for prevention and treatment of VTE and adverse pregnancy outcome. The authors acknowledged that no clear conclusions can be drawn from the studies they reviewed regarding an association between inherited thrombophilia's and adverse pregnancy outcomes-some studies show a positive

relationship, and other studies show no relationship. According to Duhl, most of the research demonstrated that FVL is not typically associated with pregnancy loss prior to 10 weeks' gestation. More evidence exists suggesting that a loss after 10 weeks' gestation may be associated with these disorders.

GeneReviews (2018): Molecular genetic testing is recommended for the following clinical scenarios:

- Individuals receiving direct thrombin inhibitors or direct factor Xa inhibitors, which may interfere with the AC resistance assay
- To confirm the diagnosis and to distinguish factor V Leiden variant heterozygotes from homozygotes and pseudohomozygotes with positive borderline or very low APC resistance assay results
- Strong lupus inhibitors and a markedly prolonged baseline aPTT when the results of testing would affect clinical management
- persons with a first unprovoked VTE who are planning to stop anticoagulation
- female relatives of persons with VTE or hereditary thrombophilia considering estrogen contraception or hormone replacement
- female relatives of persons with VTE or hereditary thrombophilia contemplating prophylactic anticoagulation during pregnancy

Outside of the US

British Committee for Standards in Hematology ([BSCH], 2010): On behalf of the BCSH, Baglin et al. noted that testing for heritable thrombophilias is not indicated in unselected patients presenting with venous thrombosis. Testing selected patients may give an indication of risk of recurrence following completion of anticoagulant therapy, for example those presenting with venous thrombosis at an early age (<40 years) and who are from apparent thrombosis-prone families (more than two other symptomatic family members). Other selected patient groups in whom the results of testing may influence treatment are children with purpura fulminans and pregnant women at risk of venous thrombosis. The decision to test these selected patients should be based on whether or not test results are likely to influence treatment decisions.

European Genetics Foundation, the Cardiovascular Disease Educational and Research Trust, the Mediterranean League on Thromboembolism, and the International Union of Angiology: In the International Consensus Statement for Thrombophilia and Thromboembolism, Nicolaidis et al. (2005) provided guidelines for investigation and management of patients with thrombophilia with or without VTE). According to the authors, heterozygote FVL is found in 0–15% of the normal population, 20% of patients with venous thrombosis and in 40% of families with thrombophilia, whereas the homozygote FVL mutation is found in only 0.02% of the normal population, but the relative risk is high. The authors researched the etiology, testing, diagnosis, risk factors, prevention and management of thrombophilia. Based on this research, the authors concluded that screening for thrombophilia should occur in:

- all patients with a first episode of spontaneous VTE
- patients with VTE under the age of 50, even with a transient predisposing factor
- patients with VTE whose only risk factor is oral contraceptive therapy or pregnancy. However, screening with other than the molecular tests should be performed at least two weeks after delivery or hormone therapy cessation.
- patients with recurrent VTE, irrespective of the presence of risk factors
- patients with recurrent superficial thrombophlebitis without cancer in the absence of varicose veins
- patients with VTE at unusual sites such as cerebral venous sinus, mesenteric or hepatic veins, and retinal vein occlusion under the age of 50
- patients with warfarin-induced skin necrosis and neonates with purpura fulminans not related to sepsis
- asymptomatic first-degree relatives of individuals with proven symptomatic thrombophilia. The authors felt this was particularly important for females in the child-bearing years.
- two consecutive or three nonconsecutive abortions at any gestational age, or one fetal death after week 20
- severe preeclampsia
- children with VTE

National Institute for Health and Care Excellence ([NICE], 2012): regarding testing for inherited thrombophilias, NICE notes the following:

- Consider testing for hereditary thrombophilia in patients who have had unprovoked DVT or PE and who have a first-degree relative who has had DVT or PE if it is planned to stop anticoagulation treatment. [2012]
- Do not offer thrombophilia testing to patients who have had provoked DVT or PE. [2012]
- Do not routinely offer thrombophilia testing to first-degree relatives of people with a history of DVT or PE and thrombophilia. [2012]

Royal College of Obstetricians and Gynaecologists (RCOG): The RCOG (2011) notes that women with second-trimester miscarriage should be tested for inherited thrombophilias including FVL, factor II (prothrombin) gene mutation, and protein S.

Amyotrophic Lateral Sclerosis:

European Federation of Neurological Societies (EFNS): Regarding amyotrophic lateral sclerosis (ALS), on behalf of the EFNS, Bergunder et al. (2011) noted that “Despite the rather low prevalence sequencing of the small SOD1 gene should be considered in patients with ALS with dominant inheritance to offer presymptomatic or prenatal diagnosis, if this is requested by the family.

Methylenetetrahydrofolate Reductase (NAD(P)H) (MTHFR):

American College of Medical Genetics (ACMG, 2013): The ACMG notes:

- MTHFR polymorphism genotyping should not be ordered as part of the clinical evaluation for thrombophilia or recurrent pregnancy loss
- MTHFR polymorphism genotyping should not be ordered for at-risk family members

Alzheimer Disease: APOE, APP, PSEN1 and PSEN2 Gene Mutations

American Academy of Neurology (AAN): The Quality Standards Subcommittee of the AAN updated an earlier practice parameter for the diagnosis of dementia in the elderly. Regarding AD, this evidence-based review concluded that there are no laboratory tests, including APOE genotyping or other genetic markers or biomarkers, which are appropriate for routine use in the clinical evaluation of patients with suspected AD. However, genotyping and biomarkers, as well as imaging, are promising avenues that are being pursued (Knopman, et al., 2001; reaffirmed 2004).

American Psychiatric Association (2007): Practice Guidelines for the treatment of patients with Alzheimer’s disease and other dementias note that a definitive diagnosis of AD requires both the clinical syndrome and microscopic examination of the brain at autopsy, at which time the characteristic plaques and neurofibrillary tangles widely distributed in the cerebral cortex will be seen. A careful clinical diagnosis of disease conforms to the pathological diagnosis 70%–90% of the time. Further, the guideline notes although genes involved in a variety of dementia syndromes have been identified and family members of patients with dementia are often concerned about their risk of developing dementia, genetic testing is generally not part of the evaluation of patients with dementia except in very specific instances. In particular, testing for apolipoprotein E4 (APOE4) is not recommended for use in diagnosis. The presence of an APOE4 allele does not change the need for a thorough workup and does not add substantially to diagnostic confidence.

National Institute on Aging (NIA, 2011): The NIA published a fact sheet noting that although a blood test can identify which APOE alleles a person has, it cannot predict who will or will not develop Alzheimer’s disease. It is unlikely that genetic testing will ever be able to predict the disease with 100% accuracy because too many other factors may influence its development and progression. Further, the NIA noted APOE testing is used in research settings to identify study participants who may have an increased risk of developing Alzheimer’s.

National Institute on Aging (NIH)/Alzheimer's Association ([AA]; 2011): The NIA/AA issued consensus recommendations regarding the diagnosis of Alzheimer's disease (AD). For probable AD dementia in a carrier of a causative genetic mutation the recommendations note that in persons who meet the core clinical criteria for probable AD dementia, evidence of a causative genetic mutation (in APP, PSEN1, or PSEN2), increases the certainty that the condition is caused by AD pathology. Carriage of the 3/4 allele of the apolipoprotein E gene is not sufficiently specific to be considered in this category.

National Society of Genetic Counselors (NSGC)/American College of Medical Genetics (ACMG): On behalf of the NSGC/ASGC, Goldman et al. (2011) published consensus practice guidelines for genetic counseling and testing for AD. The Guidelines recommend that pediatric testing for AD should not occur. Additionally, the Societies note that direct-to-consumer APOE testing is not advised.

The Guidelines note that a risk assessment should be performed by pedigree analysis to determine whether the family history is consistent with early-onset Alzheimer's disease (EOAD) or late-onset Alzheimer's disease (LOAD) and with autosomal dominant (with or without complete penetrance), familial, or sporadic inheritance. Patients should be informed that currently there are no proven pharmacologic or lifestyle choices that reduce the risk of developing AD or stop its progression. The Guidelines also note:

For families in which an autosomal dominant AD gene mutation is a possibility:

- Testing for genes associated with early-onset autosomal dominant AD should be offered in the following situations:
 - A symptomatic individual with EOAD in the setting of a family history of dementia or in the setting of an unknown family history (e.g., adoption).
 - Autosomal dominant family history of dementia with one or more cases of EOAD.
 - A relative with a mutation consistent with EOAD (currently PSEN1/2 or APP).
 - Ideally, an affected family member should be tested first. If no affected family member is available for testing and an asymptomatic individual remains interested in testing despite counseling about the low likelihood of an informative result (a positive result for a pathogenic mutation), he/she should be counseled according to the recommended protocol. If the affected relative, or their next of kin, is uninterested in pursuing testing, the option of deoxyribonucleic acid (DNA) banking should be discussed.

For families in which autosomal dominant AD is unlikely:

- Genetic testing for susceptibility loci (e.g., apolipoprotein-E [APOE]) is not clinically recommended due to limited clinical utility and poor predictive value.

Canadian Medical Association: Recommendations for risk assessment and prevention of Alzheimer's disease, based on the Third Canadian Consensus Conference on the Diagnosis and Treatment of Dementia held in March 2006, were reported by Patterson and colleagues (2008). The recommendations for genetic risk factors included:

- Predictive genetic testing may be offered to the following at-risk individuals with an apparent autosomal dominant inheritance when a family specific mutation has been identified:
 - first-degree relatives (e.g., children and siblings) of an affected person with the mutation
 - first cousins of an affected person if the common ancestors (parents who were siblings) died before the average age of onset of dementia in the family
 - nieces and nephews of an affected person whose parent (sibling of the affected person) died before the average age of onset of dementia in the family
 - minors are not usually referred for predictive genetic testing
- Genetic screening for the APOE genotype in asymptomatic individuals in the general population is not recommended because of low sensitivity and specificity.

European Federation of Neurological Societies (EFNS): On behalf of the EFNS Hort et al. (2007) published guidelines for the diagnosis and management of Alzheimer's disease. The Guideline notes "Screening for known pathogenic mutations can be undertaken in patients with appropriate phenotype or a family history of an autosomal dominant dementia. Routine apolipoprotein E (Apo E) genotyping is not recommended.

Cardiac Disease-Risk

American College of Cardiology Foundation (ACCF) /American Heart Association (AHA) Task Force on Practice Guidelines: Greenland et al. (2010) published guidelines which note that genotype testing for CHD risk assessment in asymptomatic adults is not recommended. The task force noted that there is currently no proven benefit in risk assessment when genomic testing is added to the basic global risk assessment, such as Framingham. There is no data to support results of genotype testing alter management and improve clinical outcomes.

The task force conducted a systematic review of the current scientific evidence (March 2008 – April 2010) and used evidence based methodologies to weigh the evidence which was reviewed. Level A evidence represented data from multiple randomized controlled trials or meta-analyses, level B evidence was data from a single RCT or nonrandomized trial, and level C evidence represented consensus opinion, case studies or standard of care. The recommendations were approved and endorsed by the ACCF, AHA, American Society of Echocardiography, American Society of Nuclear Cardiology, Society of Atherosclerosis Imaging and Prevention, Society of Cardiovascular Computed Tomography, and Society for Cardiovascular Magnetic Resonance. The American College of Cardiology Foundation (ACCF) and the American Heart Association (AHA) published guidelines for assessment of cardiovascular risk in asymptomatic individuals (i.e., apparently healthy adult) (Greenland, et al., 2010). The guidelines do not support genotype testing (level B evidence) or measurement of lipid parameters such as lipoproteins, apolipoproteins, particle size and density, beyond the standard fasting lipid profile (level C evidence), or natriuretic peptide testing (level B evidence).

Updated ACC/AHA guidelines on the assessment of cardiovascular risk (Goff , 2013) do not address genetic testing to determine cardiovascular risk.

Evaluation of Genomic Applications in Practice and Prevention (EGAPP, 2010): The working group concluded there was insufficient evidence to determine analytic validity, clinical validity, or clinical utility for gene expression testing for determining cardiovascular risk.

European Society of Cardiology (ESC) and the European Atherosclerosis Society (EAS): Guidelines for the management of dyslipidemias do not recommend genotype testing for cardiovascular risk estimation (Catapano, et al., 2011).

Multi-Gene Panel Testing for Non-Syndromic Hearing Loss

American College of Medical Genetics and Genomics (AMCG, 2014): Guidelines for the clinical evaluation and etiologic diagnosis of hearing loss include the following recommendations regarding genetic testing for nonsyndromic hearing loss:

- For individuals lacking physical findings suggestive of a known syndrome and having medical and birth histories that do not suggest an environmental cause of hearing loss, a tiered diagnostic approach should be implemented:
 - Pretest genetic counseling should be provided, and, with patient's informed consent, genetic testing should be ordered.
 - Single-gene testing may be warranted in cases in which the medical or family history, or presentation of the hearing loss, suggests a specific etiology.
 - In the absence of any specific clinical indications and for singleton cases and cases with apparent autosomal recessive inheritance, the next step should be testing for DFNB1-related hearing loss (due to mutations in GJB2 and adjacent deletions in GJB6). If initial genetic testing is negative, genetic testing using gene panel tests, NGS technologies such as large sequencing panels targeted toward hearing loss–related genes, whole-exome sequencing (WES), or whole-genome sequencing (WGS) strategies may be considered. Since several tests are clinically available, the clinician should be aware of the genes included in the panel chosen and the performance characteristics of the platform chosen, including coverage, analytic sensitivity, and the types of mutations that will be detected.

- If genetic testing reveals mutation(s) in a hearing loss–related gene, mutation-specific genetic counseling should be provided, followed by appropriate medical evaluations and referrals.
- If genetic testing fails to identify an etiology for a patient’s hearing loss, the possibility of a genetic or acquired etiology remains. This point should be emphasized since it may be misunderstood by clinicians and by patients and their families. For interested patients and families, further genetic testing may be pursued on a research basis.
- Regardless of whether genetic test results are positive, negative, or inconclusive, results should be communicated through the process of genetic counseling.

Coding/Billing Information

Note: 1) This list of codes may not be all-inclusive.
 2) Deleted codes and codes which are not effective at the time the service is rendered may not be eligible for reimbursement.

Laboratory Testing

Considered Not Medically Necessary:

CPT® Codes	Description
0001U	Red blood cell antigen typing, DNA, human erythrocyte antigen gene analysis of 35 antigens from 11 blood groups, utilizing whole blood, common RBC alleles reported
0084U	Red blood cell antigen typing, DNA, genotyping of 10 blood groups with phenotype prediction of 37 red blood cell antigens

Genetic Counseling

Considered Medically Necessary when criteria in the applicable policy statements listed above are met:

CPT® Codes	Description
96040	Medical genetics and genetic counseling services, each 30 minutes face-to-face with patient/family

HCPCS Codes	Description
S0265	Genetic counseling, under physician supervision, each 15 minutes

Single Gene Germline Mutation Genetic Testing

Considered Medically Necessary when criteria in the applicable policy statements listed above are met:

CPT® Codes	Description
81105	Human Platelet Antigen 1 genotyping (HPA-1), ITGB3 (integrin, beta 3 [platelet glycoprotein IIIa], antigen CD61 [GPIIIa]) (eg, neonatal alloimmune thrombocytopenia [NAIT], post-transfusion purpura), gene analysis, common variant, HPA-1a/b (L33P)
81106	Human Platelet Antigen 2 genotyping (HPA-2), GP1BA (glycoprotein Ib [platelet], alpha polypeptide [GPIba]) (eg, neonatal alloimmune thrombocytopenia [NAIT], post-transfusion purpura), gene analysis, common variant, HPA-2a/b (T145M)
81107	Human Platelet Antigen 3 genotyping (HPA-3), ITGA2B (integrin, alpha 2b [platelet glycoprotein IIb of IIb/IIIa complex], antigen CD41 [GPIIb]) (eg, neonatal alloimmune thrombocytopenia [NAIT], post-transfusion purpura), gene analysis, common variant, HPA-3a/b (I843S)
81108	Human Platelet Antigen 4 genotyping (HPA-4), ITGB3 (integrin, beta 3 [platelet glycoprotein IIIa], antigen CD61 [GPIIIa]) (eg, neonatal alloimmune thrombocytopenia [NAIT], post-transfusion purpura), gene analysis, common variant, HPA-4a/b (R143Q)

81109	Human Platelet Antigen 5 genotyping (HPA-5), ITGA2 (integrin, alpha 2 [CD49B, alpha 2 subunit of VLA-2 receptor] [GPIa]) (eg, neonatal alloimmune thrombocytopenia [NAIT], post-transfusion purpura), gene analysis, common variant (eg, HPA-5a/b (K505E))
81110	Human Platelet Antigen 6 genotyping (HPA-6w), ITGB3 (integrin, beta 3 [platelet glycoprotein IIIa, antigen CD61] [GPIIIa]) (eg, neonatal alloimmune thrombocytopenia [NAIT], post-transfusion purpura), gene analysis, common variant, HPA-6a/b (R489Q)
81111	Human Platelet Antigen 9 genotyping (HPA-9w), ITGA2B (integrin, alpha 2b [platelet glycoprotein IIb of IIb/IIIa complex, antigen CD41] [GPIIb]) (eg, neonatal alloimmune thrombocytopenia [NAIT], post-transfusion purpura), gene analysis, common variant, HPA-9a/b (V837M)
81112	Human Platelet Antigen 15 genotyping (HPA-15), CD109 (CD109 molecule) (eg, neonatal alloimmune thrombocytopenia [NAIT], post-transfusion purpura), gene analysis, common variant, HPA-15a/b (S682Y)
81161	DMD (dystrophin) (eg, Duchenne/Becker muscular dystrophy) deletion analysis, and duplication analysis, if performed.
81171	AFF2 (AF4/FMR2 family, member 2 [FMR2]) (eg, fragile X mental retardation 2 [FRAXE]) gene analysis; evaluation to detect abnormal (eg, expanded) alleles
81172	AFF2 (AF4/FMR2 family, member 2 [FMR2]) (eg, fragile X mental retardation 2 [FRAXE]) gene analysis; characterization of alleles (eg, expanded size and methylation status)
81173	AR (androgen receptor) (eg, spinal and bulbar muscular atrophy, Kennedy disease, X chromosome inactivation) gene analysis; full gene sequence
81174	AR (androgen receptor) (eg, spinal and bulbar muscular atrophy, Kennedy disease, X chromosome inactivation) gene analysis; known familial variant
81177	ATN1 (atrophin 1) (eg, dentatorubral-pallidoluysian atrophy) gene analysis, evaluation to detect abnormal (eg, expanded) alleles
81178	ATXN1 (ataxin 1) (eg, spinocerebellar ataxia) gene analysis, evaluation to detect abnormal (eg, expanded) alleles
81179	ATXN2 (ataxin 2) (eg, spinocerebellar ataxia) gene analysis, evaluation to detect abnormal (eg, expanded) alleles
81180	ATXN3 (ataxin 3) (eg, spinocerebellar ataxia, Machado-Joseph disease) gene analysis, evaluation to detect abnormal (eg, expanded) alleles
81181	ATXN7 (ataxin 7) (eg, spinocerebellar ataxia) gene analysis, evaluation to detect abnormal (eg, expanded) alleles
81182	ATXN8OS (ATXN8 opposite strand [non-protein coding]) (eg, spinocerebellar ataxia) gene analysis, evaluation to detect abnormal (eg, expanded) alleles
81183	ATXN10 (ataxin 10) (eg, spinocerebellar ataxia) gene analysis, evaluation to detect abnormal (eg, expanded) alleles
81184	CACNA1A (calcium voltage-gated channel subunit alpha1 A) (eg, spinocerebellar ataxia) gene analysis; evaluation to detect abnormal (eg, expanded) alleles
81185	CACNA1A (calcium voltage-gated channel subunit alpha1 A) (eg, spinocerebellar ataxia) gene analysis; full gene sequence
81186	CACNA1A (calcium voltage-gated channel subunit alpha1 A) (eg, spinocerebellar ataxia) gene analysis; known familial variant
81187	CNBP (CCHC-type zinc finger nucleic acid binding protein) (eg, myotonic dystrophy type 2) gene analysis, evaluation to detect abnormal (eg, expanded) alleles
81188	CSTB (cystatin B) (eg, Unverricht-Lundborg disease) gene analysis; evaluation to detect abnormal (eg, expanded) alleles
81189	CSTB (cystatin B) (eg, Unverricht-Lundborg disease) gene analysis; full gene sequence
81190	CSTB (cystatin B) (eg, Unverricht-Lundborg disease) gene analysis; known familial variant(s)
81200	ASPA (aspartoacylase) (eg, Canavan disease) gene analysis, common variants (eg, E285A, Y231X)
81204	AR (androgen receptor) (eg, spinal and bulbar muscular atrophy, Kennedy disease, X chromosome inactivation) gene analysis; characterization of alleles (eg, expanded size or methylation status)

81220	CFTR (cystic fibrosis transmembrane conductance regulator) (eg, cystic fibrosis) gene analysis; common variants (eg, ACMG/ACOG guidelines)
81221	CFTR (cystic fibrosis transmembrane conductance regulator) (eg, cystic fibrosis) gene analysis; common variants (eg, ACMG/ACOG guidelines), known familial variants
81222	CFTR (cystic fibrosis transmembrane conductance regulator) (eg, cystic fibrosis) gene analysis; duplication/deletion variants
81223	CFTR (cystic fibrosis transmembrane conductance regulator) (eg, cystic fibrosis) gene analysis; full gene sequence
81224	CFTR (cystic fibrosis transmembrane conductance regulator) (eg, cystic fibrosis) gene analysis; intron 8 poly-T analysis (eg, male infertility)
81234	DMPK (DM1 protein kinase) (eg, myotonic dystrophy type 1) gene analysis; evaluation to detect abnormal (expanded) alleles
81238	F9 (coagulation factor IX) (eg, hemophilia B), full gene sequence
81239	DMPK (DM1 protein kinase) (eg, myotonic dystrophy type 1) gene analysis; characterization of alleles (eg, expanded size)
81240 [†]	F2 (prothrombin, coagulation factor II) (eg, hereditary hypercoagulability) gene analysis, 20210G>A variant
81241 [†]	F5 (coagulation factor V) (eg, hereditary hypercoagulability) gene analysis, Leiden variant
81243	FMR1 (fragile X mental retardation 1) (eg, fragile X mental retardation) gene analysis; evaluation to detect abnormal (eg, expanded) alleles
81244	FMR1 (fragile X mental retardation 1) (eg, fragile X mental retardation) gene analysis; characterization of alleles (eg, expanded size and promoter methylation status)
81247	G6PD (glucose-6-phosphate dehydrogenase) (eg, hemolytic anemia, jaundice), gene analysis; common variant(s) (eg, A, A-)
81248	G6PD (glucose-6-phosphate dehydrogenase) (eg, hemolytic anemia, jaundice), gene analysis; known familial variant(s)
81249	G6PD (glucose-6-phosphate dehydrogenase) (eg, hemolytic anemia, jaundice), gene analysis; full gene sequence
81251	GBA (glucosidase, beta, acid) (eg, Gaucher disease) gene analysis, common variants (eg, N370S, 84GG, L444P, IVS2+1G>A)
81252	GJB2 (gap junction protein, beta 2, 26kDa; connexin 26) (eg, nonsyndromic hearing loss) gene analysis; full gene sequence
81253	GJB2 (gap junction protein, beta 2, 26kDa; connexin 26) (eg, nonsyndromic hearing loss) gene analysis; known familial variants
81254	GJB6 (gap junction protein, beta 6, 30kDa, connexin 30) (eg, nonsyndromic hearing loss) gene analysis, common variants (eg, 309kb [del(GJB6-D13S1830)] and 232kb [del(GJB6-D13S1854)])
81255	HEXA (hexosaminidase A [alpha polypeptide]) (eg, Tay-Sachs disease) gene analysis, common variants (eg, 1278insTATC, 1421+1G>C, G269S)
81256	HFE (hemochromatosis) (eg, hereditary hemochromatosis) gene analysis, common variants (eg, C282Y, H63D)
81257	HBA1/HBA2 (alpha globin 1 and alpha globin 2) (eg, alpha thalassemia, Hb Bart hydrops fetalis syndrome, HbH disease), gene analysis; common deletions or variant (eg, Southeast Asian, Thai, Filipino, Mediterranean, alpha3.7, alpha4.2, alpha20.5, and Constant Spring)
81258	HBA1/HBA2 (alpha globin 1 and alpha globin 2) (eg, alpha thalassemia, Hb Bart hydrops fetalis syndrome, HbH disease), gene analysis; known familial variant
81259	HBA1/HBA2 (alpha globin 1 and alpha globin 2) (eg, alpha thalassemia, Hb Bart hydrops fetalis syndrome, HbH disease), gene analysis; full gene sequence
81269	HBA1/HBA2 (alpha globin 1 and alpha globin 2) (eg, alpha thalassemia, Hb Bart hydrops fetalis syndrome, HbH disease), gene analysis; duplication/deletion variants
81271	HTT (huntingtin) (eg, Huntington disease) gene analysis; evaluation to detect abnormal (eg, expanded) alleles
81274	HTT (huntingtin) (eg, Huntington disease) gene analysis; characterization of alleles (eg, expanded size)

81284	FXN (frataxin) (eg, Friedreich ataxia) gene analysis; evaluation to detect abnormal (expanded) alleles
81285	FXN (frataxin) (eg, Friedreich ataxia) gene analysis; characterization of alleles (eg, expanded size)
81286	FXN (frataxin) (eg, Friedreich ataxia) gene analysis; full gene sequence
81289	FXN (frataxin) (eg, Friedreich ataxia) gene analysis; known familial variant(s)
81302	MECP2 (methyl CpG binding protein 2) (eg, Rett Syndrome) gene analysis; full sequence analysis
81303	MECP2 (methyl CpG binding protein 2) (eg, Rett Syndrome) gene analysis; known familial variant
81304	MECP2 (methyl CpG binding protein 2) (eg, Rett Syndrome) gene analysis; duplication/deletion variants
81312	PABPN1 (poly[A] binding protein nuclear 1) (eg, oculopharyngeal muscular dystrophy) gene analysis, evaluation to detect abnormal (eg, expanded) alleles
81329	SMN1 (survival of motor neuron 1, telomeric) (eg, spinal muscular atrophy) gene analysis; dosage/deletion analysis (eg, carrier testing), includes SMN2 (survival of motor neuron 2, centromeric) analysis, if performed
81330	SMPD1 (sphingomyelin phosphodiesterase 1, acid lysosomal) (eg, Niemann-Pick disease, Type A) gene analysis, common variants (eg, R496L, L302P, fsP330)
81332	SERPINA1 (serpin peptidase inhibitor, clade A, alpha-1 antiproteinase, antitrypsin, member 1) (eg, alpha-1-antitrypsin deficiency), gene analysis, common variants (eg, *S and *Z)
81333	TGFBI (transforming growth factor beta-induced) (eg, corneal dystrophy) gene analysis, common variants (eg, R124H, R124C, R124L, R555W, R555Q)
81336	SMN1 (survival of motor neuron 1, telomeric) (eg, spinal muscular atrophy) gene analysis; full gene sequence
81337	SMN1 (survival of motor neuron 1, telomeric) (eg, spinal muscular atrophy) gene analysis; known familial sequence variant(s)
81343	PPP2R2B (protein phosphatase 2 regulatory subunit Bbeta) (eg, spinocerebellar ataxia) gene analysis, evaluation to detect abnormal (eg, expanded) alleles
81344	TBP (TATA box binding protein) (eg, spinocerebellar ataxia) gene analysis, evaluation to detect abnormal (eg, expanded) alleles
81361	HBB (hemoglobin, subunit beta) (eg, sickle cell anemia, beta thalassemia, hemoglobinopathy); common variant(s) (eg, HbS, HbC, HbE)
81362	HBB (hemoglobin, subunit beta) (eg, sickle cell anemia, beta thalassemia, hemoglobinopathy); known familial variant(s)
81363	HBB (hemoglobin, subunit beta) (eg, sickle cell anemia, beta thalassemia, hemoglobinopathy); duplication/deletion variant(s)
81364	HBB (hemoglobin, subunit beta) (eg, sickle cell anemia, beta thalassemia, hemoglobinopathy); full gene sequence
81400 ^{††}	Molecular pathology procedure, Level 1 (eg, identification of single germline variant [eg, SNP] by techniques such as restriction enzyme digestion or melt curve analysis)
81401 ^{††}	Molecular pathology procedure, Level 2 (eg, 2-10 SNPs, 1 methylated variant, or 1 somatic variant [typically using nonsequencing target variant analysis], or detection of a dynamic mutation disorder/triplet repeat)
81402	Molecular pathology procedure, Level 3 (eg, >10 SNPs, 2-10 methylated variants, or 2-10 somatic variants [typically using non-sequencing target variant analysis], immunoglobulin and T-cell receptor gene rearrangements, duplication/deletion variants of 1 exon, loss of heterozygosity [LOH], uniparental disomy [UPD])
81403 ^{††}	Molecular pathology procedure, Level 4 (eg, analysis of single exon by DNA sequence analysis, analysis of > 10 amplicons using multiplex PCR in 2 or more independent reactions, mutation scanning or duplication/deletion variants of 2-5 exons)
81404 ^{††}	Molecular pathology procedure, Level 5 (eg, analysis of 2-5 exons by DNA sequence analysis, mutation scanning or duplication/deletion variants of 6-10 exons, or characterization of a dynamic mutation disorder/triplet repeat by Southern blot analysis)

81405 ^{††}	Molecular pathology procedure, Level 6 (eg, analysis of 6-10 exons by DNA sequence analysis, mutation scanning or duplication/deletion variants of 11-25 exons, regionally targeted cytogenomic array analysis)
81406 ^{††}	Molecular pathology procedure, Level 7 (eg, analysis of 11-25 exons by DNA sequence analysis, mutation scanning or duplication/deletion variants of 26-50 exons)
81407	Molecular pathology procedure, Level 8 (eg, analysis of 26-50 exons by DNA sequence analysis, mutation scanning or duplication/deletion variants of >50 exons, sequence analysis of multiple genes on one platform)
81408	Molecular pathology procedure, Level 9 (eg, analysis of > 50 exons in a single gene by DNA sequence analysis)
81479 ^{†††}	Unlisted molecular pathology procedure
83520 ^{†††}	Immunoassay for analyte other than infectious agent antibody or infectious agent antigen; quantitative, not otherwise specified

[†]Note: Considered Not Medically Necessary when used to report prothrombin gene mutation for the screening, diagnosis or management of coronary heart disease

^{††}Note: Considered Not Medically Necessary when used to report:

- ACE (angiotensin converting enzyme) (eg, hereditary blood pressure regulation), insertion/deletion variant (81400)
- AGTR1 (angiotensin II receptor, type 1) (eg, essential hypertension), 1166A>C variant (81400)
- ANG (angiogenin, ribonuclease, RNase A family, 5) (eg, amyotrophic lateral sclerosis), full gene sequence (81403)
- APOE (apolipoprotein E) (eg, hyperlipoproteinemia type III, cardiovascular disease, Alzheimer disease), common variants (eg, *2, *3, *4) (81401)
- APP (amyloid beta [A4] precursor protein) (eg, Alzheimer's disease), full gene sequence (81406)
- CDKN2A (cyclin-dependent kinase inhibitor 2A) (eg, CDKN2A-related cutaneous malignant melanoma, familial atypical mole-malignant melanoma syndrome), full gene sequence (81404)
- FUS (fused in sarcoma) (eg, amyotrophic lateral sclerosis), full gene sequence (81406)
- OPTN (optineurin) (eg, amyotrophic lateral sclerosis), full gene sequence (81406)
- PSEN1 (presenilin 1) (eg, Alzheimer's disease), full gene sequence (81405)
- PSEN2 (presenilin 2 [Alzheimer's disease 4]) (eg, Alzheimer's disease), full gene sequence (81406)
- SOD1 (superoxide dismutase 1, soluble) (eg, amyotrophic lateral sclerosis), full gene sequence (81404)
- TARDBP (TAR DNA binding protein) (eg, amyotrophic lateral sclerosis), full gene sequence (81405)

^{†††}Note: Considered Medically Necessary when used to report any covered single gene genetic test that does not have an assigned CPT/HCPCS code when criteria in the applicable policy statements listed above are not met.

HCPCS Codes	Description
S3841	Genetic testing for retinoblastoma
S3842	Genetic testing for Von Hippel-Lindau disease
S3844	DNA analysis of the connexin26 gene (GJB2) for susceptibility to congenital, profound deafness
S3845	Genetic testing for alpha-thalassemia
S3846	Genetic testing for hemoglobin E beta-thalassemia
S3849	Genetic testing for Niemann-Pick disease
S3850	Genetic testing for sickle cell anemia

S3853	Genetic testing for myotonic muscular dystrophy
-------	---

Considered Not Medically Necessary:

CPT®* Codes	Description
81291	MTHFR (5,10-methylenetetrahydrofolate reductase) (eg, hereditary hypercoagulability) gene analysis, common variants (eg, 677T, 1298C)

HCPCS Codes	Description
S3800	Genetic testing for amyotrophic lateral sclerosis (ALS)
S3852	DNA analysis for APOE epsilon 4 allele for susceptibility to Alzheimer's disease

Multi-Gene Germline Mutation Genetic Testing Panels

Considered Medically Necessary when criteria in the applicable policy statements listed above are met:

CPT®* Codes	Description
81410	Aortic dysfunction or dilation (eg, Marfan syndrome, Loeys Dietz syndrome, Ehler Danlos syndrome type IV, arterial tortuosity syndrome); genomic sequence analysis panel, must include sequencing of at least 9 genes, including FBN1, TGFBR1, TGFBR2, COL3A1, MYH11, ACTA2, SLC2A10, SMAD3, and MYLK
81430	Hearing loss (eg, nonsyndromic hearing loss, Usher syndrome, Pendred syndrome); genomic sequence analysis panel, must include sequencing of at least 60 genes, including CDH23, CLRN1, GJB2, GPR98, MTRNR1, MYO7A, MYO15A, PCDH15, OTOF, SLC26A4, TMC1, TMPRSS3, USH1C, USH1G, USH2A, and WFS1
81431	Hearing loss (eg, nonsyndromic hearing loss, Usher syndrome, Pendred syndrome); duplication/deletion analysis panel, must include copy number analyses for STRC and DFNB1 deletions in GJB2 and GJB6 genes
81442	Noonan spectrum disorders (eg, Noonan syndrome, cardio-facio-cutaneous syndrome, Costello syndrome, LEOPARD syndrome, Noonan-like syndrome), genomic sequence analysis panel, must include sequencing of at least 12 genes, including BRAF, CBL, HRAS, KRAS, MAP2K1, MAP2K2, NRAS, PTPN11, RAF1, RIT1, SHOC2, and SOS1
81448	Hereditary peripheral neuropathies (eg, Charcot-Marie-Tooth, spastic paraplegia), genomic sequence analysis panel, must include sequencing of at least 5 peripheral neuropathy-related genes (eg, BSCL2, GJB1, MFN2, MPZ, REEP1, SPAST, SPG11, SPTLC1)
81460	Whole mitochondrial genome (eg, Leigh syndrome, mitochondrial encephalomyopathy, lactic acidosis, and stroke-like episodes [MELAS], myoclonic epilepsy with ragged-red fibers [MERFF], neuropathy, ataxia, and retinitis pigmentosa [NARP], Leber hereditary optic neuropathy [LHON]), genomic sequence, must include sequence analysis of entire mitochondrial genome with heteroplasmy detection
81465	Whole mitochondrial genome large deletion analysis panel (eg, Kearns-Sayre syndrome, chronic progressive external ophthalmoplegia), including heteroplasmy detection, if performed
81479†	Unlisted molecular pathology procedure
0012U	Germline disorders, gene rearrangement detection by whole genome next-generation sequencing, DNA, whole blood, report of specific gene rearrangement(s)
0013U	Oncology (solid organ neoplasia), gene rearrangement detection by whole genome next-generation sequencing, DNA, fresh or frozen tissue or cells, report of specific gene rearrangement(s)
0014U	Hematology (hematolymphoid neoplasia), gene rearrangement detection by whole genome next-generation sequencing, DNA, whole blood or bone marrow, report of specific gene rearrangement(s)

†Note: Considered Medically Necessary when used to report any covered genetic testing panel that does not have an assigned CPT/HCPCS code

Considered Not Medically Necessary:

CPT®* Codes	Description
81440	Nuclear encoded mitochondrial genes (eg, neurologic or myopathic phenotypes), genomic sequence panel, must include analysis of at least 100 genes, including BCS1L, C10orf2, COQ2, COX10, DGUOK, MPV17, OPA1, PDSS2, POLG, POLG2, RRM2B, SCO1, SCO2, SLC25A4, SUCLA2, SUCLG1, TAZ, TK2, and TYMP
81490	Autoimmune (rheumatoid arthritis), analysis of 12 biomarkers using immunoassays, utilizing serum, prognostic algorithm reported as a disease activity score
81599††	Unlisted multianalyte assay with algorithmic analysis
82397	Chemiluminescent assay
83520††	Immunoassay for analyte other than infectious agent antibody or infectious agent antigen; quantitative, not otherwise specified
84999††	Unlisted chemistry procedure
88346	Immunofluorescence, per specimen; initial single antibody stain procedure
88350	Immunofluorescence, per specimen; each additional single antibody stain procedure (List separately in addition to code for primary procedure)
0004M	Scoliosis, DNA analysis of 53 single nucleotide polymorphisms (SNPs), using saliva, prognostic algorithm reported as a risk score

††**Note:** Considered Not Medically Necessary when used to report any non-covered genetic testing panel that does not have an assigned CPT/HCPCS code

Considered Experimental/Investigational/Unproven:

CPT®* Codes	Description
81493	Coronary artery disease, mRNA, gene expression profiling by real-time RT-PCR of 23 genes, utilizing whole peripheral blood, algorithm reported as a risk score

*Current Procedural Terminology (CPT®) ©2018 American Medical Association: Chicago, IL.

References

1. American Academy of Ophthalmology. Recommendations for genetic testing of inherited eye disease. 2014. Accessed Oct 3, 2019. Available at URL address: <http://www.aao.org/clinical-statement/recommendations-genetic-testing-of-inherited-eye-d>
2. American Academy of Pediatrics. Ethical and policy issues in genetic testing and screening in children. Pediatrics. Mar 2013. Accessed Oct 3, 2019. Available at URL address: <http://pediatrics.aappublications.org/content/131/3/620.abstract?rss=1>
3. American Board of Genetic Counseling, Inc. ABGC Certification. © 2019 American Board of Genetic Counseling, Inc. Accessed Jul 5, 2018. Available at URL Address: <http://www.abgc.net/ABGC/AmericanBoardofGeneticCounselors.asp>
4. American College of Medical Genetics and Genomics (ACMG) Board of Directors. Points to consider in the clinical application of genomic sequencing. Genet Med. 2012 Aug;14(8):759-61.
5. American College of Medical Genetics and Genomics (ACMG) Board of Directors. Points to consider for informed consent for genome/exome sequencing. Genet Med. 2013 Sep;15(9):748-9
6. American College of Medical Genetics and Genomics (ACMG). © 2001-2019 American College of Medical Genetics. Accessed Oct 3, 2019. Available at URL address: <https://www.acmg.net/>

7. American College of Obstetricians and Gynecologists. Practice Bulletin No.197: Inherited thrombophilias in pregnancy.(Jun 25, 2018). Accessed Oct 3, 2019. Available at URL address:
<https://www.acog.org/en/Clinical%20Guidance%20and%20Publications/Practice%20Bulletins%20List>
8. American Society of Human Genetics (ASHG). Genetic testing and insurance [background statement]. The Ad Hoc Committee on Genetic Testing/Insurance Issues. *Am J Hum Genet.* 1995;56:327-31.
9. American Thoracic Society; European Respiratory Society. American Thoracic Society/European Respiratory Society statement: standards for the diagnosis and management of individuals with alpha-1 antitrypsin deficiency. *Am J respire Crit Care med,* 2003 Oct 1;168(7):818-900.
10. American Urological Association (AUA).(a) The evaluation of the azoospermic male: best practice statement. Confirmed 2011. Accessed Oct 3, 2019. Available at URL address:
<http://www.auanet.org/content/guidelines-and-quality-care/clinical-guidelines.cfm>
11. American Urological Association (AUA).(b) The management of obstructive azoospermia: AUA best practice statement. Confirmed 2011. Accessed Oct 3, 2019. Available at URL address:
<https://www.auanet.org/search?keywords=azoospermia>
12. American Urological Association (AUA). (c) The optimal evaluation of the infertile male: AUA best practice statement. Confirmed 2011. Accessed Oct 3, 2019. Available at URL address:
[http://www.auanet.org/guidelines/male-infertility-optimal-evaluation-\(reviewed-and-validity-confirmed-2011\)](http://www.auanet.org/guidelines/male-infertility-optimal-evaluation-(reviewed-and-validity-confirmed-2011))
13. Anderson JL1, Knight S, May HT, Horne BD, Bair TL, Huntinghouse JA, Rollo JS, Muhlestein JB, Carlquist JF. Validation and quantification of genetic determinants of lipoprotein-a levels and predictive value for angiographic coronary artery disease. *Am J Cardiol.* 2013 Sep 15;112(6):799-804.
14. Anderson PM, Borasio GD, Dengler R, Hardiman O, Kollwe K, Leigh PN, et al. EFNS task force on management of amyotrophic lateral sclerosis: guidelines for diagnosing and clinical care of patients and relatives. An evidence-based review with good practice points. *Eur J Neurol.* 2005 Dec;12(12): 921-38.
15. Ashley-Koch A, Yang Q, Olney RS. Sickle hemoglobin (HbS) allele and sickle cell disease: a HuGE review. *Am J Epidemiol.* 2000 May 1;151(9):839-45.
16. Baglin T, Gray E, Greaves M, Hunt BJ, Keeling D, Machin S, et al. Clinical guidelines for testing for heritable thrombophilia. *Br J Haematol.* 2010 Apr;149(2):209–20. Accessed Jul 5, 2018. Available at URL address: <https://onlinelibrary.wiley.com/doi/epdf/10.1111/j.1365-2141.2009.08022.x>
17. Bare LA, Morrison AC, Rowland CM, Shiffman D, Luke MM, Iakoubova OA, Kane JP, Malloy MJ, Ellis SG, Pankow JS, Willerson JT, Devlin JJ, Boerwinkle E. Five common gene variants identify elevated genetic risk for coronary heart disease . *Genet Med.* 2007 Oct;9(10):682-9.
18. Bates SM, Greer IA, Middeldorp S, Veenstra DL, Prabulos AM, Vandvik PO, American College of Chest Physicians. VTE, thrombophilia, antithrombotic therapy,and pregnancy: Antithrombotic Therapy and Prevention of Thrombosis, 9th ed:American College of Chest Physicians Evidence-Based Clinical Practice Guidelines.*Chest.* 2012 Feb;141(2 Suppl):e691S-736S.
19. Benderly M, Boyko V, Goldbourt U. Apolipoproteins and long-term prognosis in coronary heart disease patients. *Am Heart J.* 2009 Jan;157(1):103-10.
20. Bermudez EA, Rifai N, Buring J, Manson JE, Ridker PM. Interrelationships among circulating interleukin-6, C-reactive protein, and traditional cardiovascular risk factors in women. *Arterioscler Thromb Vasc Biol.* 2002 Oct 1;22(10):1668-73.

21. Bird TD. Alzheimer disease overview. GeneReviews®. Initial Posting: October 23, 1998; Last Revision: September 24, 2015. Accessed Oct 3, 2019. Available at URL address: <https://www.ncbi.nlm.nih.gov/books/NBK1161/>
22. Bird TD. Early-onset familial Alzheimer disease. GeneReview®. Initial Posting: September 24, 1999; Last Revision: October 18, 2012. Accessed Oct 3, 2019. Available at URL address: <https://www.ncbi.nlm.nih.gov/books/NBK1236/>
23. Biron-Andreani C, Bauters A, Le Cam-Duchez V, Delahousse B, Lequerrec A, Dutrillaux F, et al. Factor v leiden homozygous genotype and pregnancy outcomes. *Obstetrics & Gynecology*. 2009 Dec;114(6):1249-53.
24. Biswas S, Ghoshal PK, Mandal SC, Mandal N. Association of low-density lipoprotein particle size and ratio of different lipoproteins and apolipoproteins with coronary heart disease. *J Cardiol*. 2008 Oct;52(2):118-26.
25. Bradley LA, Palomaki GE, Bienstock J, Varga E, Scott JA. Can factor V Leiden and prothrombin G20210A testing in women with recurrent pregnancy loss result in improved pregnancy outcomes? Results from a targeted evidence-based review. *Genet Med*. 2012 Jan;14(1):39-50. Epub 2011 Sep 13.
26. Brunklaus A, Dorris L, Ellis R, Reavey E, Lee E, Forbes G, Appleton R, Cross JH, Ferrie C, Hughes I, Jollands A, King MD, Livingston J, Lynch B, Philip S, Scheffer IE, Williams R, Zuberi SM, et al. The clinical utility of an SCN1A genetic diagnosis in infantile-onset epilepsy. *Dev Med Child Neurol*. 2013 Feb;55(2):154-61.
27. Burgunder JM, Schols L, Baets J, Anderson P, Gasser T, Szolnoki Z, et al. EFNS guidelines for the molecular diagnosis of neurogenetic disorders: motoneuron, peripheral nerve and muscle disorders. *Eur J Neurol*. 2011 Feb;18(2):207-17.
28. Burke W. Genetic testing. *N Engl J Med*. 2002 Dec 5;347(23):1867-75.
29. Bushnell C, McCullough LD, Awad IA, et al; American Heart Association Stroke Council; Council on Cardiovascular and Stroke Nursing; Council on Clinical Cardiology; Council on Epidemiology and Prevention; Council for High Blood Pressure Research. Guidelines for the prevention of stroke in women: a statement for healthcare professionals from the American Heart Association/American Stroke Association. *Stroke*. 2014 May;45(5):1545-88.
30. Caga-anan ECF, Smith L, Sharp RR, Lantos JD. Testing children for adult-onset genetic diseases. *Pediatrics*. 2012 Jan;129(1):163-7.
31. Cantürk C, Baade U, Salazar R, Storm N, Pörtner R, Höppner W. Sequence analysis of CYP21A1P in a German population to aid in the molecular biological diagnosis of congenital adrenal hyperplasia. *Clin Chem*. 2011 Mar;57(3):511-7.
32. Catapano AL, Reiner Z, De Backer G, Graham I, Taskinen MR, Wiklund O, et al. ESC/EAS Guidelines for the management of dyslipidaemias The Task Force for the management of dyslipidaemias of the European Society of Cardiology (ESC) and the European Atherosclerosis Society (EAS). *Atherosclerosis*. 2011 Jul;217(1):3-46.
33. Chasman DI, Shiffman D, Zee RY, Louie JZ, Luke MM, Rowland CM, Catanese JJ, Buring JE, Devlin JJ, Ridker PM. Polymorphism in the apolipoprotein(a) gene, plasma lipoprotein(a), cardiovascular disease, and low-dose aspirin therapy. *Atherosclerosis*. 2009 Apr;203(2):371-6.
34. Chew EY, Klein ML, Clemons TE, Agrón E, Abecasis GR. Genetic Testing in Persons with Age-Related Macular Degeneration and the Use of the AREDS Supplements: To Test or Not to Test? *Ophthalmology*. 2015 Jan;122(1):212-5.

35. Chinnery P, Majamaa K, Turnbull D, Thorburn D. Treatment for mitochondrial disorders. *Cochrane Database Syst Rev*. 2006 Jan 25;(1):CD004426.
36. Chinnery PF, Turnbull DM. Clinical features, investigation, and management of patients with defects of mitochondrial DNA. *J Neurol Neurosurg Psychiatry*. 1997 Nov;63(5):559-63.
37. Chong LY, Fenu E, Stansby G, Hodgkinson S. Management of venous thromboembolic diseases and the role of thrombophilia testing: summary of NICE guidance. *BMJ*. 2012 Jun 27;344:e3979.
38. Clarke, R, Peden, JF, Hopewell, JC, et al. Genetic variants associated with Lp(a) lipoprotein level and coronary disease. *N Engl J Med*. 2009 Dec 24;361(26):2518-28.
39. Crews KR, Gaedigk A, Dunnenberger HM, et al. Clinical pharmacogenetics implementation consortium guidelines for cytochrome P450 2D6 genotype and codeine therapy: 2014 update. *Clin Pharmacol Ther*. 2014 Apr;95(4):376-82. Epub 2014 Jan 23.
40. Croles FN, Nasserinejad K, Duvekot JJ, Kruip MJ, Meijer K, Leebeek FW. Pregnancy, thrombophilia, and the risk of a first venous thrombosis: systematic review and Bayesian meta-analysis. *BMJ*. 2017 Oct 26;359:j4452.
41. Culling B, Ogle R. Genetic counseling issues in cystic fibrosis. *Paediatr Respir Rev*. 2010 Jun;11(2):75-9.
42. Daniels SE, Beineke P, Rhees B, McPherson JA, Kraus WE, Thomas GS, Rosenberg S. Biological and analytical stability of a peripheral blood gene expression score for obstructive coronary artery disease in the PREDICT and COMPASS studies. *J Cardiovasc Transl Res*. 2014 Oct;7(7):615-22. Epub 2014 Aug 14.
43. de Vries MC, Rodenburg RJ, Morava E et al. Normal biochemical analysis of the oxidative phosphorylation (OXPHOS) system in a child with POLG mutations: a cautionary note. *J Inherit Metab Dis*. 2008 Dec;31 Suppl 2:S299-302.
44. Donley G, Hull SC, Berkman BE. Prenatal whole genome sequencing: just because we can, should we? *Hastings Cent Rep*. 2012 Jul-Aug;42(4):28-40.
45. Duncan RD, Prucka S, Wiatrak BJ, Smith RJ, Robin NH. Pediatric otolaryngologists' use of genetic testing. *Arch Otolaryngol Head Neck Surg*. 2007 Mar;133(3):231-6.
46. Egan M, Green DM, Voynow JA. Cystic fibrosis. In: Kliegman RM, Stanton BF, St. Geme III, JW, Schor NF, editors *Nelson Textbook of Pediatrics*, 20th ed. St. Louis: WB Saunders: 2016.
47. Elashoff MR, Wingrove JA, Beineke P, et al. Development of a Blood-Based Gene Expression Algorithm for Assessment of Obstructive Coronary Artery Disease in Non-Diabetic Patients. *BMC Med Genomics* 2011; 4(1):26doi: 10.1186/1755-8794-4-26.
48. Elliott et al. Pathogenic mitochondrial DNA mutations are common in the general population. *Am J Hum Genet*. 2008 Aug;83(2):254-60.
49. Evaluation of Genomic Applications in Practice and Prevention (EGAPP) Working Group. Recommendations from the EGAPP Working Group: genomic profiling to assess cardiovascular risk to improve cardiovascular health. *Genet Med* 2010 Dec;12(12):839-43.
50. Evaluation of Genomic Applications in Practice and Prevention (EGAPP) Working Group. Recommendations from the EGAPP Working Group: routine testing for Factor V Leiden (R506Q) and prothrombin (20210G>A) mutations in adults with a history of idiopathic venous thromboembolism and their adult family members. *Genet Med*. 2011 Jan;13(1):67-76.

51. Farbstein D, Levy AP. The genetics of vascular complications in diabetes mellitus. *Cardiol Clin of North America*. Aug 2010;28(3).
52. Filsoof DM, Safford RE, Newby K, Rosenberg S, Kontras DG, Baker A, Odunukan OW, Fletcher G. Impact of exercise stress testing on diagnostic gene expression in patients with obstructive and nonobstructive coronary artery disease. *Am J Cardiol*. 2015 May 15;115(10):1346-50. doi: 10.1016/j.amjcard.2015.02.041. Epub 2015 Feb 18.
53. Finucane B, Abrams L, Cronister A, Archibald AD, Bennett RL, McConkie-Rosell A. Genetic counseling and testing for FMR1 gene mutations: practice guidelines of the National Society of Genetic Counselors. *J Genet Couns*. 2012 Dec;21(6):752-60.
54. Folsom AR, Nambi V, Pankow JS, Tang W, Farbaksh K, Yamagishi K, Boerwinkle E. Effect of 9p21 genetic variation on coronary heart disease is not modified by other risk markers. The Atherosclerosis Risk in Communities (ARIC) Study. *Atherosclerosis*. 2012 Oct;224(2):435-9.
55. Gardner P, Oitmaa E, Messner A, Hoefsloot L, Metspalu A, Schrijver I. Simultaneous multigene mutation detection in patients with sensorineural hearing loss through a novel diagnostic microarray: a new approach for newborn screening follow-up. *Pediatrics*. 2006 Sep;118(3):985-94.
56. Geerts WH, Bergqvist D, Pineo GF, Heit JA, Samama CM, Lassen MR, Colwell CW; American College of Chest Physicians. Prevention of venous thromboembolism: American College of Chest Physicians Evidence-Based Clinical Practice Guidelines (8th Edition). *Chest*. 2008 Jun;133(6 Suppl):381S-453S.
57. Genetic Testing Registry. National Center for Biotechnology Information, U.S. National Library of Medicine. Accessed Oct 3, 2019. Available at URL address: <http://www.ncbi.nlm.nih.gov/gtr/>
58. GenoMEL. Genetics Melanoma Consortium. CDKN2A. Referral guidelines for genetic counseling and testing. Oct 2011. Accessed Oct 3, 2019. Available at URL address: http://www.genomel.org/genetic_counselling/
59. Glanz K, Volpicelli K, Kanetsky PA, Ming ME, Schuchter LM, Jepson C, Domchek SM, Armstrong K. Melanoma genetic testing, counseling, and adherence to skin cancer prevention and detection behaviors. *Cancer Epidemiol Biomarkers Prev*. 2013 Apr;22(4):607-14.
60. Goff DC Jr, Lloyd-Jones DM, Bennett G, Coady S, D'Agostino RB, Gibbons R, et al. 2013 ACC/AHA guideline on the assessment of cardiovascular risk: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines. *Circulation*. 2014 Jun 24;129(25 Suppl 2):S49-73. doi: 10.1161/01.cir.0000437741.48606.98. Epub 2013 Nov 12. Erratum in: *Circulation*. 2014 Jun 24;129(25 Suppl 2):S74-5. PubMed PMID: 24222018. Accessed Oct 3, 2019. Available at URL address: https://www.clinicalkey.com/service/content/pdf/watermarked/1-s2.0-S0735109713060312.pdf?locale=en_US
61. Graham BH. Diagnostic challenges of mitochondrial disorders: complexities of two genomes. *Methods Mol Biol*. 2012;837:35-46.
62. Green RC, Berg JS, Grody WW, Kalia SS, Korf BR, Martin CL, McGuire AL, Nussbaum RL, O'Daniel JM, Ormond KE, Rehm HL, Watson MS, Williams MS, Biesecker LG. ACMG recommendations for reporting of incidental findings in clinical exome and genome sequencing. *Genet Med*. 2013 Jul;15(7):565-74.
63. Greenland P, Alpert JS, Beller GA, Benjamin EJ, Budoff MJ, Fayad ZA, et al. 2010 ACCF/AHA guideline for assessment of cardiovascular risk in asymptomatic adults: a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines. *J Am Coll Cardiol*. 2010 Dec 14;56(25):e50-103.

64. Gregg AR, Simpson JL. Genetic screening for cystic fibrosis. *Obstet Gynecol Clin North Am.* 2002 Jun; 29(2): 329-40.
65. Grody W, Cutting GR, Klinger KW, Richards CS, Watson MS, Desnick RJ (Subcommittee on Cystic Fibrosis Screening, Accreditation of Genetic Services Committee, ACMG). Laboratory standards and guidelines for population-based cystic fibrosis carrier screening. *Genet Med.* 2001 Mar/Apr;3(2):149-54. Updated 2008. Accessed Jul 5, 2018. Available at URL address: <https://www.acmg.net/PDFLibrary/Cystic-Fibrosis-Population-Based-Carrier-Screening-Standards.pdf>
66. Grody W, Griffin JH, Taylor AK, Korf BR, Heit JA. American College of Medical Genetics consensus statement on Factor Leiden mutation testing. *Genet Med.* 2001 Mar-Apr;3(2):139-48.
67. Grody WW, Thompson BH, Gregg AR, Bean LH, Monaghan KG, Schneider A, Lebo RV. ACMG position statement on prenatal/preconception expanded carrier screening. *Genet Med.* 2013 Jun;15(6):482-3.
68. Gross SJ, Pletcher BA, Monaghan KG; Professional Practice and Guidelines Committee. Carrier screening in individuals of Ashkenazi Jewish descent. *Genet Med.* 2008 Jan;10(1):54-6.
69. Hansen CB, Wadge LM, Lowstuter K, Boucher K, Leachman SA. Clinical germline genetic testing for melanoma. *Lancet Onc.* 2004;5(5).
70. Herman L, Froelich J, Kanelos D, et al. Utility of a genomic-based, personalized medicine test in patients presenting with symptoms suggesting coronary artery disease. *J Am Board Fam Med.* 2014;27(2):258-267.
71. Hickey SE, Curry CJ, Toriello HV. ACMG Practice Guideline: lack of evidence for MTHFR polymorphism testing. *Genet Med.* 2013 Feb;15(2):153-6.
72. Holtzman NE, Watson MS, editors. Final Report of the Task Force on Genetic Testing: promoting safe and effective genetic testing in the United States. 1997 Sep. Last reviewed Oct 1, 2012. Accessed Jul 2018. Available at URL address: <http://www.genome.gov/page.cfm?pageID=10001733>
73. Hone SW, Smith RJ. Genetic screening for hearing loss. *Clin Otolaryngol Allied Sci.* 2003 Aug;28(4):285-90.
74. Hopewell JC1, Clarke R, Parish S, Armitage J, Lathrop M, Hager J, Collins R; Heart Protection Study Collaborative Group. Lipoprotein(a) genetic variants associated with coronary and peripheral vascular disease but not with stroke risk in the Heart Protection Study. *Circ Cardiovasc Genet.* 2011 Feb;4(1):68-73.
75. Hresko MT. Clinical practice. Idiopathic scoliosis in adolescents. *N Engl J Med.* 2013; 368(9):834-841.
76. Human Genome Project Information. Modified Apr 23, 2019. Accessed Oct 3, 2019. Available at URL address: http://www.ornl.gov/sci/techresources/Human_Genome/home.shtml
77. Iakoubova OA, Sabatine MS, Rowland CM, Tong CH, Catanese JJ, Ranade K, Simonsen KL, Kirchgessner TG, Cannon CP, Devlin JJ, Braunwald E. Polymorphism in KIF6 gene and benefit from statins after acute coronary syndromes: results from the PROVE IT-TIMI 22 study. *J Am Coll Cardiol.* 2008 Jan 29;51(4):449-55.
78. Jensen EH, Borja-Cacho D, Al-Refaie WB, Vickers SM. Exocrine pancreas. In: Townsend CM, Beauchamp RD, Evers BM, Mattox KL, editors. *Sabiston textbook of surgery. The biological basis of modern surgical practice*, 20th Elsevier, Philadelphia (PA), 2017.
79. Johansen Taber KA, Dickinson BD, Wilson M. The promise and challenges of next-generation genome sequencing for clinical care. *JAMA Intern Med.* 2014 Feb 1;174(2):275-80.

80. Johnson AD, Newton-Cheh C, Chasman DI, Ehret GB, Johnson T, Rose L, et al. Association of hypertension drug target genes with blood pressure and hypertension in 86,588 individuals. *Hypertension*. 2011 May;57(5):903-10.
81. Kamien BA, Cardamone M, Lawson JA, Sachdev R A genetic diagnostic approach to infantile epileptic encephalopathies. *J Clin Neurosci*. 2012 Jul;19(7):934-41. Epub 2012 May 20.
82. Kelly E, Greene CM, Carroll TP, McElvaney NG, O'Neill SJ. Alpha-1 antitrypsin deficiency. *Respir Med*. 2010 Jun;104(6):763-72. doi: 10.1016/j.rmed.2010.01.016. Epub 2010 Mar 20.
83. Keren A, Syrris P, McKenna WJ. Hypertrophic cardiomyopathy: the genetic determinants of clinical disease expression. *Nat Clin Pract Cardiovasc Med*. 2008 Mar;5(3):158-68.
84. Khoury MJ, McCabe LL, McCabe ERB. Population screening in the age of genomic medicine. *N Engl J Med*. 2003 Jan 3;348(1):50-8.
85. Koch, W, Mueller, JC, Schrempf, M, et al. Two rare variants explain association with acute myocardial infarction in an extended genomic region including the apolipoprotein(A) gene. *Annals of human genetics*. 2013 Jan;77(1):47-55.
86. Krintus M, Kozinski M, Kubica J, Sypniewska G. Critical appraisal of inflammatory markers in cardiovascular risk stratification. *Crit Rev Clin Lab Sci*. 2014 Oct;51(5):263-79. doi: 10.3109/10408363.2014.913549. Epub 2014 Jun 11.
87. Ladapo JA(1), Herman L, Weiner BH, Rhees B, Castle L, Monane M, McPherson JA. Use of a blood test incorporating age, sex, and gene expression influences medical decision-making in the evaluation of women presenting with symptoms suggestive of obstructive coronary artery disease: summary results from two ambulatory care studies in primary care. *Menopause*. 2015 Apr 6. [Epub ahead of print]
88. Ladapo JA, Lyons H, Yau M, et al. Enhanced assessment of chest pain and related symptoms in the primary care setting through the use of a novel personalized medicine genomic test: Results from a prospective registry study. *Am J Med Qual*. 2014 May 5.
89. Lansky A, Elashoff MR, Ng V, et al. A gender-specific blood-based gene expression score for assessing obstructive coronary artery disease in nondiabetic patients: results of the Personalized Risk Evaluation and Diagnosis in the Coronary Tree (PREDICT) trial. *Am Heart J*. 2012 Sep;164(3):320-6.
90. Li ZG1, Li G, Zhou YL, Chen ZJ, Yang JQ, Zhang Y, Sun S, Zhong SL. Lack of association between lipoprotein(a) genetic variants and subsequent cardiovascular events in Chinese Han patients with coronary artery disease after percutaneous coronary intervention. *Lipids Health Dis*. 2013 Aug 27;12:127.
91. Lieb W, Pavlik R, Erdmann J, Mayer B, Holmer SR, Fischer M, Baessler A, Hengstenberg C, Loewel H, Doering A, Riegger GA, Schunkert H. No association of interleukin-6 gene polymorphism (-174 G/C) with myocardial infarction or traditional cardiovascular risk factors. *Int J Cardiol*. 2004 Nov;97(2):205-12.
92. Lieber DS, Calvo SE, Shanahan K, et al. Targeted exome sequencing of suspected mitochondrial disorders. *Neurology*. 2013;80(19):1762–70.
93. MacCallum P, Bowles L, Keeling D. Diagnosis and management of heritable thrombophilias. *BMJ*. 2014 Jul 17;349:g4387.
94. March of Dimes. Newborn Screening Tests. March of Dimes Birth Defect Foundation. © 2019 March of Dimes Foundation. Accessed Oct 3, 2019. Available at URL address: http://www.marchofdimes.com/professionals/14332_1200.asp

95. Marchiori A, Mosena L, Prins MH, Prandoni P. The risk of recurrent venous thromboembolism among heterozygous carriers of factor V Leiden or prothrombin G20210A mutation. A systematic review of prospective studies. *Haematologica*. 2007 Aug;92(8):1107-14.
96. Marciniul DD, Hernandez P, Balter M, Bourbeau J, Chapman KR, Ford GT, et al. Canadian Thoracic Society COPD Clinical Assembly Alpha-1 Antitrypsin Deficiency Expert Working Group. Alpha-1 antitrypsin deficiency targeted testing and augmentation therapy: a Canadian Thoracic Society clinical practice guideline. *Can Respir J*. 2012 Mar-Apr;19(2):109-16. Eratum in: *Can Respir J*. 2012 Jul-Aug;19(4):272.
97. Marlin S, Feldmann D, Blons H, Loundon N, Rouillon I, Albert S, et al. GJB2 and GJB6 mutations: genotypic and phenotypic correlations in a large cohort of hearing-impaired patients. *Arch Otolaryngol Head Neck Surg*. 2005 Jun;131(6):481-7.
98. Maron BJ. Hypertrophic cardiomyopathy. In: Mann, DL, Zipes DP, Libby P, Bonow RO, editors. *Braunwald's heart disease: a textbook of cardiovascular medicine*, 11th ed. Elsevier, Philadelphia (PA), 2019.
99. McCormick E1, Place E, Falk MJ. Molecular genetic testing for mitochondrial disease: from one generation to the next. *Neurotherapeutics*. 2013 Apr;10(2):251-61.
100. McPherson JA, Davis K, et al. The clinical utility of gene expression testing on the diagnostic evaluation of patients presenting to the cardiologist with symptoms of suspected obstructive coronary artery disease: results from the IMPACT (Investigation of a Molecular Personalized Coronary Gene Expression Test on Cardiology Practice Pattern) trial. *Crit Path Cardiol*. 2013 Jun;12(2):37-42.
101. Michelson DJ, Shevell MI, Sherr EH, Moeschler JB, Gropman AL, Ashwal S. Evidence report: Genetic and metabolic testing on children with global developmental delay: report of the Quality Standards Subcommittee of the American Academy of Neurology and the Practice Committee of the Child Neurology Society. *Neurology*. 2011 Oct 25;77(17):1629-35.
102. Murray B, Mitsumoto H. Disorders of upper and lower motor neurons. In: Daroff RB, Jankovic J, Mazziotti JC, Pomeroy SL, editors. *Bradley's Neurology in clinical practice*. 7th edition. Elsevier, Inc. Philadelphia (PA); 2016.
103. National Comprehensive Cancer Network® (NCCN) NCCN GUIDELINES™ Clinical Guidelines in Oncology™. © National Comprehensive Cancer Network, Inc. 2019, All Rights Reserved. Clinical practice guidelines in oncology. Accessed Oct 3, 2019. Available at URL address: <https://www.nccn.org>
104. National Health Service (NHS). National Educational and Development Center. Genetics education. Accessed Oct 3, 2019. Available at URL address: <http://www.geneticseducation.nhs.uk/>
105. National Institute for Health and Clinical Excellence. © 2019 National Institute for Health and Care Excellence. Accessed Oct 3, 2019. Available at <http://www.nice.org.uk/Guidance>.
106. National Center for Biotechnology Information (NCBI). GeneReviews. Accessed Oct 3, 2019. Available at URL address: <http://www.ncbi.nlm.nih.gov/books/>
107. National Institutes of Health (NIH). Genetics Home Reference. Accessed Oct 3, 2019. Available at URL address: <https://ghr.nlm.nih.gov/>
108. National Institutes of Health. National Institute of Neurological Disorders and Stroke (NINDS). Amyotrophic lateral sclerosis fact sheet. Updated 2019 Mar 27. Accessed Oct 3, 2019. Available at URL address: <https://www.ninds.nih.gov/Disorders/All-Disorders/Amyotrophic-Lateral-Sclerosis-ALS-Information-Page>
109. National Reference Center for Bioethics Literature. The Joseph and Rose Kennedy Institute of Ethics, Georgetown University. Scope note 22. Genetic testing and genetic screening. Sept 1993. Updated Jul

2011. Accessed Oct 3, 2019. Available at URL address:
<https://repository.library.georgetown.edu/bitstream/handle/10822/556877/sn22.pdf?sequence=1>

110. National Society of Genetic Counselors. Position statement on incidental findings in genetic testing. Updated Apr 16, 2019. Accessed Oct 3, 2019. Available at URL address:
<https://www.nsgc.org/p/bl/et/blogaid=30>
111. Nauck M, Winkelmann BR, Hoffmann MM, Böhm BO, Wieland H, März W. The interleukin-6 G(-174)C promoter polymorphism in the LURIC cohort: no association with plasma interleukin-6, coronary artery disease, and myocardial infarction. *J Mol Med*. 2002 Aug;80(8):507-13.
112. Nimkarn S, New MI. Congenital adrenal hyperplasia due to 21-hydroxylase deficiency: A paradigm for prenatal diagnosis and treatment. *Ann N Y Acad Sci*. 2010 Mar;1192:5-11.
113. Ogura Y, Takahashi Y, Kou I, et al. A replication study for association of 53 single nucleotide polymorphisms in a scoliosis prognostic test with progression of adolescent idiopathic scoliosis in Japanese. *Spine (Phila Pa 1976)*. 2013; 38(16):1375-1379.
114. Pagliarini DJ, Calvo SE, Chang BA et al. A mitochondrial protein compendium elucidates complex I disease biology. *Cell*. 2008 Jul 11;134(1):112-23. doi: 10.1016/j.cell.2008.06.016
115. Palomaki GE, Melillo S, Bradley LA. Association between 9p21 genomic markers and heart disease: a meta-analysis. *JAMA*. 2010a Feb 17;303(7):648-56.
116. Pfeffer G, Horvath R, Klopstock T et al. New treatments for mitochondrial disease-no time to drop our standards. *Nat Rev Neurol*. 2013 Aug;9(8):474-81. doi: 10.1038/nrneurol.2013.129.
117. Practice Committee of the American Society for Reproductive Medicine. Electronic address: ASRM@asrm.org; Practice Committee of the American Society for Reproductive Medicine. Combined hormonal contraception and the risk of venous thromboembolism: a guideline. *Fertil Steril*. 2017 Jan;107(1):43-51.
118. Prior TW. Carrier screening for spinal muscular atrophy. *Genet Med*. 2008 Nov. 10(11):840-2.
119. Prior TW, Narasimhan N, Sugarman EA, Batish SD, Braastad C. Technical standards and guidelines for spinal muscular atrophy testing. *Genet Med*. 2011 Jul;13(7):686-94.
120. Priori SG, Napolitano C, Gasparini M. Natural history of Brugada syndrome. Insights for risk stratification and management. *Circulation*. 2002;105:1342.
121. Pruthi RK. Optimal utilization of thrombophilia testing. *Int J Lab Hematol*. 2017 May;39 Suppl 1:104-110.
122. Putcha GV, Bejjani BA, Bleoo S, Booker JK, Carey JC, Carson N, et al. A multicenter study of the frequency and distribution of GJB2 and GJB6 mutations in a large North American cohort. *Genet Med*. 2007 Jul;9(7):413-26. Pezzolesi MG, Krolewski AS. The genetic risk of kidney disease in type 2 diabetes. *Med Clin North America*. Jan 2013;97(1).
123. Raffini LJ, Scott JP. Hereditary predisposition to thrombosis. In: Kleigman RM, Stanton BF, St. Geme III, JW, Schor NF, editors. *Nelsons textbook of pediatrics*, 20th ed. Elsevier. 2016
124. Ratnapriya R, Chew EY. Age-related macular degeneration-clinical review and genetics update. *Clin Genet*. 2013 Aug;84(2):160-6.
125. Reichert S, Schlitt A, Benten AC, Hofmann B, Schaller HG, Schulz S. The interleukin 6 c.-174 CC genotype is a predictor for new cardiovascular events in patients with coronary heart disease within three years follow-up. *Cytokine*. 2016 Jul;83:136-138.

126. Ridker PM, Libby P, Buring JE. Risk factors for atherothrombotic disease. In: Mann, DL, Zipes DP, Libby P, Bonow RO, editors. Braunwald's heart disease: a textbook of cardiovascular medicine, 11th ed. Elsevier, Philadelphia (PA), 2019.
127. Rodgers MA, Betancourt MT, Clark P, Lindqvist PZ, Dizon-Townson D, Said J, et al. The association of factor V Leiden and prothrombin gene mutation and placenta-mediated pregnancy complications: a systematic review and meta-analysis of prospective cohort studies.
128. Rosenberg S, Elashoff MR, Lieu HD, et al., PREDICT Investigators. Whole blood gene expression testing for coronary artery disease in nondiabetic patients: major adverse cardiovascular events and interventions in the PREDICT trial. *J Cardiovasc Transl Res.* 2012 Jun;5(3):366-74.
129. Rosenberg S, Elashoff MR, Beineke P, et al. Multicenter validation of the diagnostic accuracy of a blood-based gene expression test for assessing obstructive coronary artery disease in nondiabetic patients. *Ann Intern Med.* 2010 Oct 5;153(7):425-34.
130. Rühle F, Stoll M. Advances in predicting venous thromboembolism risk in children. *Br J Haematol.* 2018 Mar;180(5):654-665.
131. Sacconi S, Salviati L, Sue CM et al. Mutation screening in patients with isolated cytochrome c oxidase deficiency. *Pediatr Res.* 2003 Feb;53(2):224-30.
132. Sarafoglou K, Lorentz CP, Otten N, Oetting WS, Grebe SK. Molecular testing in congenital adrenal hyperplasia due to 21 α -hydroxylase deficiency in the era of newborn screening. *Clin Genet.* 2012 Jul;82(1):64-70.
133. Savage SK, Ziniel SI, Stoler J, Marguiles DM, Holm IA, Brownstein CA. An assessment of clinician and researcher needs for support in the era of genomic medicine. *Personalized Medicine.* 2014; 11(6): 569-579. doi: 10.2217/pme.14.48.
134. Scott DA, Yoshinaga-Itano C, ACMG Working Group on Update of Genetics Evaluation Guidelines for the Etiologic Diagnosis of Congenital Hearing Loss: Professional Practice and Guidelines Committee. *Genet Med.* 2014 Apr;16(4): 347-55. Epub 2014 Mar 20.
135. Segal JB, Brotman DJ, Necochea AJ, Emadi A, Samal L, Wilson LM, et al. Predictive value of factor V Leiden and prothrombin G20210A in adults with venous thromboembolism and in family members of those with a mutation: a systematic review. *JAMA.* 2009 Jun 17;301(23):2472-85.
136. Shashi V, McConkie-Rosell A, Rosell B, Schoch K, Vellore K, McDonald M, Jiang YH, Xie P, Need A, Goldstein DB. The utility of the traditional medical genetics diagnostic evaluation in the context of next-generation sequencing for undiagnosed genetic disorders. *Genet Med.* 2014 Feb;16(2):176-82.
137. Shaw PJ. Amyotrophic lateral sclerosis and other motor neuron disease. In: Goldman L, Schafer AI, editors. *Goldman's Cecil medicine*, 25th ed. Elsevier Saunders; Philadelphia (PA), 2016.
138. Shearer AE, DeLuca AP, Hildebrand MS, Taylor KR, Gurrola J 2nd, Scherer S, et al. Comprehensive genetic testing for hereditary hearing loss using massively parallel sequencing. *Proc Natl Acad Sci U S A.* 2010 Dec 7;107(49):21104-9.
139. Shearer AE, Kolbe DL, Azaiez H, Sloan CM, Frees KL, Weaver AE, et al. Copy number variants are a common cause of non-syndromic hearing loss. *Genome Med.* 2014 May 22;6(5):37.
140. Siemering K, Manji SS, Hutchison WM, Du Sart D, Phelan D, Dahl HH. Detection of mutations in genes associated with hearing loss using a microarray-based approach. *J Mol Diagn.* 2006 Sep;8(4):483-9.

141. Shiffman D, Chasman DI, Zee RY, Iakoubova OA, Louie JZ, Devlin JJ, Ridker PM. A kinesin family member 6 variant is associated with coronary heart disease in the Women's Health Study. *J Am Coll Cardiol*. 2008a Jan 29;51(4):444-8.
142. Shiffman D, O'Meara ES, Bare LA, et al. Association of gene variants with incident myocardial infarction in the Cardiovascular Health Study. *Arterioscler Thromb Vasc Biol*. 2008b Jan;28(1):173-9.
143. Sie MP, Sayed-Tabatabaei FA, Oei HH, et al. Interleukin 6 -174 g/c promoter polymorphism and risk of coronary heart disease: results from the rotterdam study and a meta-analysis. *Arterioscler Thromb Vasc Biol*. 2006 Jan;26(1):212-7.
144. Snoeckx RL, Huygen PL, Feldmann D, Marlin S, Denoyelle F, Waligora J, et al. GJB2 mutations and degree of hearing loss: a multicenter study. *Am J Hum Genet*. 2005 Dec;77(6):945-57.
145. Stevens SM, Woller SC, Bauer KA, Kasthuri R, Cushman M, Streiff M, et al. Guidance for the evaluation and treatment of hereditary and acquired thrombophilia. *J Thromb Thrombolysis*. 2016 Jan;41(1):154-64.
146. Sukhija R, Fahdi I, Garza L, Fink L, Scott M, Aude W, Pacheco R, Bursac Z, Grant A, Mehta JL. Inflammatory markers, angiographic severity of coronary artery disease, and patient outcome. *Am J Cardiol*. 2007 Apr 1;99(7):879-84.
147. Tang QL, Julien C, Eveleigh R, et al. A Replication Study for Association of 53 Single Nucleotide Polymorphisms in ScolioScore™ Test with Adolescent Idiopathic Scoliosis in French-Canadian Population. *Spine (Phila Pa 1976)*. 2015 Feb 2
148. Tester DJ, Ackerman MJ. Genetics of cardiac arrhythmias. In: Mann DL, Zipes DP, Libby P, Bonow RO, Braunwald E, editors. *Braunwald's heart disease: a textbook of cardiovascular medicine*, 11th ed. Elsevier, Philadelphia (PA); 2019.
149. Thomas GS, Voros S, McPherson JA, Lansky AJ, Winn ME, Bateman TM, Elashoff MR, Lieu HD, Johnson AM, Daniels SE, Ladapo JA, Phelps CE, Douglas PS, Rosenberg S. A blood-based gene expression test for obstructive coronary artery disease tested in symptomatic nondiabetic patients referred for myocardial perfusion imaging the COMPASS study. *Circ Cardiovasc Genet*. 2013 Apr;6(2):154-62.
150. Tsai MY, Loria CM, Cao J, Kim Y, Siskovick D, Schreiner PJ, et al. Clinical utility of genotyping the 677C>T variant methylenetetrahydrofolate reductase in humans is decreased in the post folic acid fortification era. *J Nutr*. 2009 Jan;139(1):33-7.
151. U.S. Preventive Services Task Force. Screening for Idiopathic Scoliosis in Adolescents. November 2004. AHRQ Pub. No. 05-0568-A
152. Van Driest SL, Jaeger MA, Ommen SR, Will ML, Gersh BJ, Tajik AJ, et al. Comprehensive analysis of the beta-myosin heavy chain gene in 389 unrelated patients with hypertrophic cardiomyopathy. *J Am Coll Cardiol*. 2004 Aug 4;44(3):602-10.
153. Van Driest SL, Ommen SR, Tajik AJ, Gersh BJ, Ackerman MJ. Yield of genetic testing in hypertrophic cardiomyopathy. *Mayo Clin Proc*. 2005 Jun;80(6):739-44.
154. van El CG, Cornel MC, Borry P, Hastings RJ, Fellmann F, Hodgson SV, Howard HC, Cambon-Thomsen A, Knoppers BM, Meijers-Heijboer H, Scheffer H, Tranebjaerg L, Dondorp W, de Wert GM; ESHG Public and Professional Policy Committee. Whole-genome sequencing in health care: recommendations of the European Society of Human Genetics. *Eur J Hum Genet*. 2013 Jun;21(6):580-4.
155. van Ommen CH, Nowak-Göttl U. Inherited Thrombophilia in Pediatric Venous Thromboembolic Disease: Why and Who to Test. *Front Pediatr*. 2017 Mar 14;5:50.

156. Vargas J, Lima JA, Kraus WE, Douglas PS, Rosenberg S. use of the Corus CAD Gene Expression Test for assessment of obstructive coronary artery disease likelihood in symptomatic non-diabetic patients. *PLoS Curr*. 2013 Aug 26;5.
157. Vassy JL, Meigs JB. Is genetic testing useful to predict type 2 diabetes? *Best Pract Res Clin Endocrinol Metab*. 2012 Apr;26(2):189-201. doi: 10.1016/j.beem.2011.09.002.
158. Waldemar G, Dubois B, Emre M, Georges J, McKeith IG, Rossor M et al. EFNS. Recommendations for the diagnosis and management of Alzheimer's disease and other disorders associated with dementia: EFNS guideline. *Eur J Neurol*. 2007 Jan;14(1):e1-26.
159. Ward K, Lazovich D, Hordinsky MK. Germline melanoma susceptibility and prognostic genes: a review of the literature. *J Am Acad Dermatol*. 2012;67:1055-67.
160. Ward K, Ogilvie JW, Singleton MV, Chettier R, Engler G, Nelson LM. Validation of DNA-based prognostic testing to predict spinal curve progression in adolescent idiopathic scoliosis. *Spine (Phila Pa 1976)*. 2010; 35(25):E1455-1464.
161. Wijesekera LC, Leigh PN. Amyotrophic lateral sclerosis. *Orphanet J Rare Dis*. 2009 Feb 3;4:3.
162. Winand R, Hens K, Dondorp W, de Wert G, Moreau Y, Vermeesch JR, Liebaers I, Aerts J. In vitro screening of embryos by whole-genome sequencing: now, in the future or never? *Hum Reprod*. 2014 Feb 2.
163. Wingrove JA, Daniels SE, Sehnert AJ et al. Correlation of peripheral-blood gene expression with the extent of coronary artery stenosis. *Circ Cardiovasc Genet* 2008; 1(1):31-8.
164. Wolf N, Smeitink JA. Mitochondrial disorders: a [proposal for consensus diagnostic criteria in infants and children. *Neurology*. 2002; Nov12;59(9): 1402-5.
165. Wong LJ(1), Naviaux RK, Brunetti-Pierri N, Zhang Q, Schmitt ES, Truong C. Molecular and clinical genetics of mitochondrial diseases due to POLG mutations. *Hum Mutat*. 2008 Sep;29(9):E150-72. doi: 10.1002/humu.20824.
166. Wong LJ, Scaglia F, Graham BH et al. Current molecular diagnostic algorithm for mitochondrial disorders. *Mol Genet Metab*. 2010 Jun;100(2):111-7. doi: 10.1016/j.ymgme.2010.02.024.
167. Wu O, Robertson L, Twaddle S, Lowe GDO, Clark P, Greaves M, Walker ID, et al. Screening for thrombophilia in high-risk situations: systematic review and cost-effectiveness analysis. The Thrombosis: Risk and Economic Assessment of Thrombophilia Screening (TREATS) study. *Health Technol Assess*. 2006 Apr;10(11):1-110.
168. Yang Y, Muzny DM, Reid JG, Bainbridge MN, Willis A, Ward PA, et al. Clinical whole-exome sequencing for the diagnosis of mendelian disorders. *N Engl J Med*. 2013 Oct 17;369(16):1502-11
169. Yang XR, Pfeiffer RM, Wheeler W et al. Identification of modifier genes for cutaneous malignant melanoma in melanoma-prone families with and without CDKN2A mutations. *Int J Cancer* 25(12):2912-7.
170. Wong LJ, Scaglia F, Graham BH et al. Current molecular diagnostic algorithm for mitochondrial disorders. *Mol Genet Metab*. 2010 Jun;100(2):111-7. doi: 10.1016/j.ymgme.2010.02.024.
171. Zawati MH, Parry D, Thorogood A, Nguyen MT, Boycott KM, Rosenblatt D, Knoppers BM. Reporting results from whole-genome and whole-exome sequencing in clinical practice: a proposal for Canada? *J Med Genet*. 2014 Jan;51(1):68-70.

172. Zeller T, Blankenberg S. Blood-based gene expression tests: Promises and limitations. *Circ Cardiovasc Genet.* 2013;6(2):139-140.

"Cigna Companies" refers to operating subsidiaries of Cigna Corporation. All products and services are provided exclusively by or through such operating subsidiaries, including Cigna Health and Life Insurance Company, Connecticut General Life Insurance Company, Cigna Behavioral Health, Inc., Cigna Health Management, Inc., QualCare, Inc., and HMO or service company subsidiaries of Cigna Health Corporation. The Cigna name, logo, and other Cigna marks are owned by Cigna Intellectual Property, Inc. © 2019 Cigna.